

Recommendations

Neurosurgical management of the acute phase of adult and pediatric traumatic brain injury: 2025 guidelines of the French Society of Neurosurgery



ARTICLE INFO

Keywords:

Brain injuries
Traumatic
Craniocerebral trauma
Intracranial pressure
Skull fractures
Neurosurgical procedures
Practice guidelines
Hydrocephalus
Critical care

ABSTRACT

Objective: To develop a multidisciplinary French framework addressing neurosurgical management in the initial phase of traumatic brain injury (TBI) in adults and children.

Design: A panel of 29 experts was formed at the request of the French Society of Neurosurgery (SFNC), with the participation of the French Society of Pediatric Neurosurgery (SFNCP), French Society of Private-Practice Neurosurgeons (SFNCL), French-Speaking Neurocritical Care and Neuro-Anesthesiology Society (ANARLF), French Society of Anesthesia, Critical Care and Perioperative Medicine (SFAR), French-Speaking Pediatric Emergency and Intensive Care Group (GFRUP), French Society of Neuroradiology (SFNR), French-Speaking Infectious Diseases Society (SPILF), and the French Society of Physical Medicine and Rehabilitation (SOFMER).

Methods: Questions were formulated using the PICO (Patients, Intervention, Comparison, Outcome) format, grouped into 7 categories: 1. Factors of poor prognosis, 2. Extradural hematoma, 3. Acute subdural hematoma, 4. Skull-base fracture and dural tear, 5. Penetrating traumatic brain injury, 6. Post-traumatic cerebrospinal fluid disorder, and 7. Pediatric specificities.

Results: Synthesis by the experts and application of the GRADE® method resulted in the formulation of 45 recommendations. Strong consensus was reached for all recommendations at the first round of rating.

Conclusion: There was a strong consensus among the experts on important interdisciplinary recommendations to improve the neurosurgical management of patients with TBI.

1. Introduction

Traumatic brain injury (TBI) is one of the most serious types of trauma, accounting for 30% of post-traumatic mortality in the United States [1] and 37% in Europe [2]. Annual incidence in Europe is approximately 250/100,000, with significant regional differences [3]. Despite major efforts in pre-hospital and in-hospital management, even mild TBI remains a source of disabling neurological sequelae.

Numerous national and international clinical guidelines and consensus statements have been developed to help practitioners adopt common decision-making rules in this area. Most were developed by professional societies in critical care and emergency medicine in collaboration with experts in neurosurgery, neuroradiology and rehabilitation. There are recommendations for the medical management of mild TBI (Glasgow Coma Score ≥ 13) and concussion [4,5], for severe (GCS ≤ 8) and moderate (GCS 9–12) TBI in adults [6,7] and for TBI in children [8], as well as recommendations for pre-hospital management [9] and for intracranial brain monitoring [10,11]. In addition, the Neurocritical Care Society issued recommendations on neuroprognosis in moderate and severe TBI [12].

Although neurosurgical procedures are an integral part of this management, the only guidelines developed specifically by neurosurgeons date back to 2006 [13]. Some surgical procedures were included

in these recommendations, and not included in the present guidelines: external ventricular drainage (EVD) in adults [6,7] and children [8], and decompressive craniectomy in adults [6,7,14] and children [8]. However, the indications and modalities of neurosurgical treatment, regardless of the severity of TBI, for both adults and children are a broad area of debate. This is the subject of the present recommendations.

2. Methods

These guidelines were drawn up by a group of experts for the French Society of Neurosurgery (SFNC), in partnership with the French Society of Paediatric Neurosurgery (SFNCP), French Society of Private-Practice Neurosurgeons (SFNCL), French-Speaking Neurocritical Care and Neuro-Anesthesiology Society (ANARLF), French Society of Anesthesia, Critical Care and Perioperative Medicine (SFAR), French-Speaking Pediatric Emergency and Intensive Care Group (GFRUP), French Society of Neuroradiology (SFNR), French-Speaking Infectious Diseases Society (SPILF) and French Society of Physical Medicine and Rehabilitation (SOFMER).

The organizing committee compiled a list of questions to be addressed and designated experts for each question. Questions were formulated using the PICO (Patient Intervention Comparison Outcome) format.

<https://doi.org/10.1016/j.neuchi.2025.101686>

Available online 23 May 2025

0028-3770/© 2025 The Authors. Published by Elsevier Masson SAS. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

The GRADE® method was used to develop the guidelines. Following a quantitative analysis of the literature, the method determined the quality of evidence: i.e., the estimated confidence in the effect, and the level of recommendation. Quality of evidence was graded as:

- high: further research is very unlikely to change the confidence in the estimate of the effect;
- moderate: further research is likely to affect the confidence in the estimate of the effect and may change the estimate of the effect itself;
- low: further research is very likely to affect the confidence in the estimate of the effect and may change the estimate of the effect itself;
- very low: any estimate of the effect is very unlikely.

There are a large number of studies of neurosurgical management of head trauma in its early stages, but very few with a high level of evidence; thus, the “recommendations for professional practice” format is appropriate, allowing experts to make suggestions based on “expert opinion”, even with very weak evidence. No level 1 recommendations were made. Recommendations were written using the following standard formulations: “It is probably advisable” for Grade 2 recommendations and “The experts suggest” for expert opinion. Each recommendation was then assessed by each of the experts and rated on a scale from 1 (strongest disagreement) to 9 (strongest agreement).

For a recommendation to be made, at least 50% of voters had to have an opinion, with fewer than 20% voting for the opposite proposal. A strong consensus required agreement by at least 70% of voters.

3. Results

The scope of these guidelines is the neurosurgical management of adult and pediatric TBI requiring neurosurgical intervention in the initial phase (days after trauma). A bibliographic search was performed using the PubMed and Cochrane databases. To be included in the analysis, publications had to be published after 2006, which is the date of the previous neurosurgical guidelines, or earlier if deemed important by the expert panel, in peer-reviewed journals, in English or French. A specific analysis of the pediatric literature was performed.

The experts selected 15 questions, covering areas related to the indications and modalities of neurosurgical procedures in the treatment of early-phase TBI: prognostic factors, extradural hematoma, acute subdural hematoma, depressed skull fracture and skull-base fracture associated with cerebrospinal fluid (CSF) fistula, penetrating TBI, post-traumatic CSF disorders, and pediatric specificities. Poor neurological outcome on the Glasgow Outcome Scale Extended (GOSE) [15] was used as the primary endpoint: death (GOSE 1) and severe neurological disability (GOSE 2–4) (see Appendix). For pediatric questions, GOSE was replaced by the Pediatric Cerebral Performance Category (PCPC) in some publications [16]; PCPC > 3 was considered pejorative. Central nervous system infections and epilepsy also counted as poor neurological outcomes.

After synthesizing the work of the experts and applying the GRADE methodology, 45 recommendations were formulated. These represent a collective effort on the part of practitioners to homogenize practice and provide arguments to support neurosurgical decision-making in individual cases, despite low levels of evidence in the literature.

The recommendations were summarized in two infographics (See Appendix).

1. Factors for poor prognosis when emergency neurosurgery is considered in adult patients with severe traumatic brain injury

Question 1: In patients with severe TBI for whom emergency neurosurgery is considered, should the decision to withhold or withdraw surgical treatment be discussed in a collegial multidisciplinary setting?

In neurosurgical emergencies, early neuroprognostication within the first 72 h is difficult. The surgical decision may have far-reaching and irreversible consequences. Given the risk of severe postoperative

disability, there are situations of devastating brain injury in which abstention from surgery is indicated, withholding or withdrawing life-sustaining treatment: early life-sustaining treatment, within the first 72 h, is the leading cause of death after severe TBI, accounting for 45%–87% of in-hospital mortality [17]. Moreover, decompressive craniectomy protects the patient from death from intracranial hypertension, but at the potential cost of severe neurological disability [18]. However, the clinical and paraclinical variables to establish this prognosis at admission are limited: initial clinical examination may be carried out under uncertain conditions, the patient is not available to the neurosurgeon in case of telemedicine, and brain CT can be misleading if performed too early or may not fully explore the brainstem. Prognostic models for TBI do not accurately predict neurological outcome, explaining only 35% of variation in final prognosis [19]. A combination of a large number of modalities is necessary to refine neuro-prediction [20], and this is not feasible in very early phases.

In severe TBI, the patient is comatose and cannot take an active part in therapeutic decision-making. However, what constitutes poor prognosis is not only strictly medical and, may be judged differently by physicians, patients and relatives [21].

In practice, the surgical management of severe TBI is heterogeneous [17,22].

The framework for limitation and withdrawal of life-sustaining treatment in France and existing recommendations on prognosis and therapeutic limitations in patients with severe cerebral palsy are worth bearing in mind here.

Firstly, limitation and withdrawal of life-sustaining treatment are defined in French law: the “Kouchner law” (Act 200 2-3 03 of March 4, 2002), “Leonetti law” (Act 2005-370 of April 22, 2005) and “Claeys-Leonetti law” (Act 2016-87 of February 2, 2016) [23]. In emergency settings, strict application of the law can be difficult to enforce, and, in 2018, the French Society of Emergency Medicine (SFMU) and the French-Speaking Society of Intensive Care (SRLF) published recommendations, available online, on limitation and withdrawal of life-sustaining treatment in adults in emergency settings [24].

Secondly, the Neurocritical Care Society issued guidelines on the management of “devastating brain injury” [25], defined as any neurological condition on admission immediately threatening vital and functional prognosis and for which limitation of early intervention is envisaged [26]:

- *We recommend determining prognosis from repeated examinations over time to establish greater confidence and accuracy (strong recommendation, moderate quality of evidence).*
- *We recommend applying these guidelines in the early stages of devastating brain injury in order to maintain physiologic stability, even when early limitation of aggressive care is being considered. Such early implementation prevents unwarranted deterioration and allows sufficient opportunity for prognostic evaluation, care planning, and consideration of organ donation (strong recommendation, moderate quality of evidence).*
- *We recommend that clinicians consider all known prognostic variables in determining risk of death and that prognostication be based on individualized assessment of risk factors rather than on clinical scoring systems (strong recommendation, moderate quality of evidence).*

Recommendations

- **R 1.1:** The experts suggest that, in patients with severe traumatic brain injury who require emergency neurosurgery (within the first 72 h) but present features of a devastating brain injury, any decision to withhold neurosurgical intervention should result from a collegial discussion involving, whenever possible, three senior physicians: a neurosurgeon, a specialist in anesthesiology and critical care (ideally with expertise in neurocritical care), and, preferably, a rehabilitation physician.

EXPERT OPINION (STRONG AGREEMENT)

- **R 1.2: The experts suggest implementing a 'limited collegial decision-making process' during the period of permanent care, involving at least two senior physicians, including a neurosurgeon.**

EXPERT OPINION (STRONG AGREEMENT)

Question 2: In patients with severe TBI requiring emergency neurosurgery, is frailty associated with poor prognosis?

Frailty corresponds to age-related decline in multi-organ function, which is more or less rapid and intense depending on the individual, and better reflects the individual's vulnerability to stressors than age considered in isolation [27]. Several indices have recently been developed to assess frailty: the modified Frailty Index (mFI-11 [28]), and the shorter mFI-5 [29]), shown to be equally reliable in trauma patients [30]; the Clinical Frailty Scale (CFS) [31]; the Autonomy Maintenance Service Integration Research Program (*Programme de Recherche sur l'Intégration des Services de Maintien de l'Autonomie*: PRISMA 7) questionnaire [32]; and the Trauma-Specific Frailty Index (TSFI) [33]. These indices were shown to correlate with short- and medium-term mortality, complications, hospital stay, post-hospital functional decline and quality of life in many pathological contexts [34]. Frailty is a factor for poor prognosis independently of age in trauma patients [35].

These indices have rarely been applied in TBI: our literature review found only 3 non-randomized prospective studies [36–38], 10 retrospective studies [39–48] and 1 meta-analysis [49]. Several of these studies showed significant associations between age, frailty and neurological prognosis in TBI patients [36,39–41,49]. The CENTER TBI Frailty Index [36] is the only TBI-specific index. Although very accurate and well suited for research, its complexity precludes use in emergency settings, whereas mFI-5 ≥ 2 [37] and CFS ≥ 4 [48] are easy to determine and are reliable markers of pathological frailty in TBI patients.

Recommendations

- **R 2.1: In patients with severe TBI requiring emergency neurosurgery, it is probably advisable to consider frailty in neurosurgical decision-making.**

GRADE 2 (STRONG AGREEMENT)

- **R 2.2: The experts suggest that the following frailty scores should be considered pathological, in order to assess prognosis:**
 - **modified Frailty Index-5 (mFI-5) ≥ 2**
 - **Clinical Frailty Scale (CFS) ≥ 4**

EXPERT OPINION (STRONG AGREEMENT)

Question 3: In patients with severe TBI requiring emergency neurosurgery, is bilateral unreactive mydriasis associated with poor prognosis?

The literature reports overall poor prognosis, particularly in the absence of surgical intervention [50–59]. Most studies were subject to numerous biases, including the risk of self-fulfilling prophecy, as bilateral unreactive mydriasis is traditionally considered to have no hope of a favorable neurological outcome. Other causes of pupil dilation should be borne in mind: beta2-mimetics (salbutamol), levodopa, anticholinergic drugs (atropine, scopolamine, nefopam, serotonin reuptake inhibitors, imipramine), sympathomimetic drugs (adrenaline), central nervous system stimulants (amphetamines, cocaine), barbiturates, benzodiazepines, severe hypothermia, trauma to the eyeball and/or oculomotor nerves, etc. A recent study highlighted overestimation of mortality and of poor prognosis (GOSE ≤ 4) in the IMPACT and CRASH studies and the need to recalibrate these prognostic scores [60].

One meta-analysis [61] included 5 retrospective cohort studies for a total of 82 patients with bilateral unreactive pupils. Analysis showed favorable outcome (GOSE > 4) in 54.3% (95% CI 36.3%–71.8%) of

patients with extradural hematoma and 6.6% (95% CI 1.8%–14.1%) of patients with acute subdural hematoma.

A systematic review of the literature including 22 studies and 503 patients, in mainly low-quality studies [62], suggested that patients presenting transtentorial herniation and bilateral unreactive mydriasis at initial management who underwent ultra-early (≤ 2 h) decompression craniectomy or craniotomy had approximately a 1-in-3 chance of survival and a 1-in-6 chance of achieving functional independence (GOSE > 4). In a few small retrospective studies, aggressive management combining osmotherapy and neurosurgical treatment within 2 h of trauma was associated with favorable neurological outcome (GOSE > 4) in 26.8–44.4 % of patients [63,64].

Recommendations

- **R 3: In patients with severe TBI requiring emergency neurosurgery, the experts suggest that bilateral unreactive mydriasis (ideally assessed on automated pupillometry) should be considered as a factor for poor prognosis, but should not, in isolation, contraindicate the procedure, particularly if the mydriasis lasts for less than 2 h.**

EXPERT OPINION (STRONG AGREEMENT)

Question 4: In patients with severe TBI who require emergency neurosurgery, is the interval to surgery associated with prognosis?

The literature reports divergent results regarding the association between time to surgery and prognosis in TBI patients.

Most studies [65–78] reported a negative impact of longer intervals, while

many retrospective studies with small numbers of patients found no association [79–90], as corroborated by other studies [91–96], questioning the concept of a 'golden hour' in TBI management [97].

Recommendations

- **R 4: In patients with severe TBI who require emergency neurosurgery, the experts suggest that time to treatment should not be considered in isolation when assessing prognosis and making neurosurgical decisions.**

EXPERT OPINION (STRONG AGREEMENT)

Question 5: In patients with severe TBI requiring emergency neurosurgery, is preoperative antithrombotic treatment associated with poor prognosis?

The head injury population, which is increasingly older, is often treated with antithrombotic (AT) agents, including antiplatelet agents (APA) and anticoagulants (AC), including direct oral anticoagulants (DOA).

In some studies, patients previously treated with dual APA [98] or a combination of AC and APA [99] had a more severe clinical and radiological presentation at admission after traumatic brain injury (TBI) than controls.

Overall, APA monotherapy is said not appear to affect prognosis in TBI. Many studies were imprecise and did not distinguish between the different APAs, and it seems that this statement concerns aspirin only. Other APAs, and especially clopidogrel, are probably associated with higher risk of intracranial bleeding.

A systematic review of the literature, including 2,447 TBI patients treated with APAs and 4,814 untreated TBI patients, found no difference in the mortality between the two groups [100]. Retrospective studies showed that outcome in patients treated with APAs or ACs who had neurosurgery for TBI did not differ compared to patients without ATs [101–103].

However, several studies found increased risk of mortality in TBI patients treated with ACs. In a retrospective study of 771 patients aged ≥ 65 years, use of AT agents was associated with a risk of lesions worsening and death (58% vs 26.7%, $p < 0.05$), particularly in the group with cerebral contusion [104]. In another retrospective cohort study of

1,365 patients aged ≥ 65 years, patients receiving ATs ($n = 724$) had more intracerebral hematomas ($p < 0.0001$), worse neurological prognosis ($\text{GOSE} \leq 4$; $p < 0.0001$) and higher mortality ($p < 0.0001$). This increased mortality was found particularly in patients on dual APAs (odds ratio [OR], 4.66; 95% confidence interval [CI], 1.57–13.87), warfarin (OR 5.18; 2.15–12.51) and DOAs (OR 5.09; 1.37–18.87), but not in patients on APA mono-therapy [105]. In a third study, including 3,031 adult TBI patients, use of APAs (4% of patients) or ACs (11%) prior to TBI was independently associated with excess mortality (OR 1.62, 95% CI 1.02–2.58; and OR 1.43, 95% CI 1.06–1.94, respectively) [106]. In the subgroup aged ≥ 70 years, excess mortality at 1 year was even higher (OR 2.28, 95% CI 1.16–4.22; and OR 1.50, 95% CI 0.97–2.32, respectively). A retrospective analysis of a prospective cohort of 20,303 TBI patients aged ≥ 65 years also found a higher risk of mortality in those previously treated with ACs (OR = 1.306, $p < 0.001$) [107]. Similarly, in a retrospective cohort study of 8,312 patients aged ≥ 65 years, including 4,680 on ATs, risk of mortality or poor prognosis was higher in those treated with warfarin (OR 1.60; 95% CI 1.27–2.01), AC + APAs (OR 1.61; 95% CI 1.18–2.21) or DOAs (OR 1.67; 95% CI 1.07–2.59) compared with patients without ACs [108]. Finally, in a recent systematic review of the literature including 4,417 patients from 12 studies, prior treatment with ACs was associated with increased risk of mortality (OR 2.39; 95% CI 1.63–3.50) [109].

Use of ACs or APA + ACs prior to TBI was associated with an increased risk of poor neurological outcome ($\text{GOSE} \leq 4$) in survivors [107,110–113], with increased risk of postoperative intracranial bleeding (OR 12.242; $p = 0.0070$) [114].

Recommendations

- **R 5.1:** In patients with severe TBI who require emergency neurosurgery, particularly those over 65 years of age, it is probably advisable to consider the presence of anticoagulant treatment as an independent factor of poor outcome.

GRADE 2 (STRONG AGREEMENT)

- **R 5.2:** In patients with severe TBI requiring emergency neurosurgery, it is probably not advisable to consider the presence of anti-platelet monotherapy with aspirin as a poor prognostic factor.

GRADE 2 (STRONG AGREEMENT)

2: EXTRA-DURAL HEMATOMA

Cyrille CAPEL, Jean Denis MOYER

Question 6: Is surgical evacuation of extra-dural hematoma associated with improved neurological outcome in TBI patients?

Initial Glasgow Coma Score (GCS) is one of the most important factors in surgical decision-making and prognosis [115–119]. A decrease of at least 2 points in GCS between admission and surgery is associated with poor prognosis (GOS (Glasgow Outcome Score) < 4) [118].

The Brain Trauma Foundation guidelines [116] suggest that > 30 mL extradural hematoma (EDH) should be managed surgically, regardless of clinical status. Several studies found that patients with EDH > 25 – 30 mL were at greater risk of neurological deterioration [115, 119–123].

Frontotemporal and temporal locations are at greater risk of volume increase. Delayed surgical management was reported to be a factor for poor prognosis (22.7% GOS < 4) [115].

The swirl sign corresponds to hypodensity within the hyperdensity of the EDH, indicating active bleeding. It is present in 14–30% of EDHs [124,125] and is associated with poor prognostic signs (pupil dilation, impaired vigilance, large EDH volume) [124–126]. Mean GOS was 4.43 ± 1.25 in patients without swirl sign, compared to 3.58 ± 1.77 for those with [125].

For EDH located in posterior cerebral fossa, the Brain Trauma Foundation recommends surgery for volume > 10 mL, thickness > 15 mm

or midline deviation > 5 mm [127]. In cases of ventricular dilatation, external ventricular drainage or endoscopic ventriculocisternostomy should be considered in addition to surgical evacuation of the EDH. In these conditions, good outcome (GOS 4–5) was found in a majority of patients [128,129]. The best outcomes were in patients with a GCS ≥ 9 at the time of surgery.

EDH of venous origin (dural sinus wounds) can be difficult to evacuate and is usually managed conservatively, except in cases of large volume or neurological signs (impaired alertness, focal signs). The literature on this topic is limited. A review of the literature [130] and a retrospective study [131] found good prognosis (GOS 5) in operated patients.

With conservative treatment, the risk of poor outcome is higher if the CT scan is performed too early. In a 2002 analysis [132], trauma-to-scan time was shorter (1.3 vs. 2.3 h) in patients with volume change than in those without. Risk of poor outcome (GOS < 4) was greater if the scan was performed less than 6 h after trauma [118]. In a 2015 retrospective study of 797 patients with EDH, 98.1% of EDHs reached their final size within 5–6 h of trauma. If possible, it is therefore recommended to perform CT scan 6 h after TBI [116,118,133,134]. If the patient's condition deteriorates (impaired vigilance or focal neurological signs), the scan should be performed earlier, to guide surgical indications [118, 119]. Special attention should also be paid to conservatively treated EDH when a contralateral lesion was previously operated on, to prevent rapid worsening after surgical decompression [135].

Recommendations

- **R 6.1:** The experts suggest that the emergency surgical evacuation of extradural hematoma can improve neurological outcome in TBI patients in the following circumstances:
 - GCS ≤ 8 ;
 - Pupil dilation;
 - Estimated EDH volume > 30 mL;
 - Midline deviation > 5 mm;
 - Brainstem compression;
 - Sign of active bleeding (swirl sign).

EXPERT OPINION (STRONG AGREEMENT)

- **R 6.2:** The experts suggest that conservative treatment of EDH of arterial origin should be considered in TBI patients in the absence of clinical or radiologic severity.

EXPERT OPINION (STRONG AGREEMENT)

- **R 6.3:** The experts suggest that individualized management of EDH of venous origin should be considered in TBI patients.

EXPERT OPINION (STRONG AGREEMENT)

- **R 6.4:** The experts suggest that follow-up of patients with non-operated EDH should be performed in a center with neurosurgical expertise.

EXPERT OPINION (STRONG AGREEMENT)

- **R 6.5:** In patients with post-traumatic EDH treated conservatively, it is probably advisable to perform systematic CT 6 h post-trauma if the initial CT scan was too early, even in the absence of subsequent clinical deterioration.

GRADE 2 (STRONG AGREEMENT)

3: ACUTE SUBDURAL HEMATOMA

Matthieu FAILLOT, Stéphanie SIGAUT, Alice ROLLAND

Question 7: Is the surgical evacuation of acute subdural hematoma associated with improved neurological outcome in TBI

patients?

The Brain Trauma Foundation recommends evacuation of any acute subdural hematoma (ASDH) associated with a Glasgow Coma Score (GCS) ≤ 8 and/or with thickness >10 mm and/or with midline shift >5 mm [136]. These criteria were based on retrospective series [137–139] with young patients. Furthermore, the recommendations did not take account of negative criteria such as age, low GCS or the presence of bilateral unreactive pupils (see section 1). Overall, the management of patients with traumatic ASDH is heterogeneous [140].

In elderly patients, the risk-benefit ratio is difficult to establish. Early evacuation of SDH in these patients was associated with significant morbidity [141–144]. However, initial conservative management (which may be followed by burr-hole evacuation) might lead to significant neurological and/or systemic deterioration associated with prolonged hospital stay [145–147]. The biases of retrospective studies in older patients prevented demonstration of clear benefit of surgery in this population [142,143,145–147], as management was very heterogeneous [148]. The only randomized trial that might have addressed this issue [149] was halted due to recruitment difficulties [150]. In this context, there is no justification for withholding neurosurgical treatment in older patients with low frailty scores, particularly in the absence of associated intraparenchymal lesions. Regarding initial neurological status, patients operated on for ASDH with initial GCS of 3 showed very high mortality, at 65–100% [61,151,152]. Regarding patients operated for ASDH with bilaterally unreactive pupils, a systematic review found 66% mortality, and favorable neurological outcome in only 7% of patients [61].

In practice, studies found a trend for lower mortality with surgery, but no improvement in neurological outcome [153]. A meta-analysis found lower mortality in operated comatose patients (GCS ≤ 8) with an absolute risk reduction of 23%–40% [154]. Two recent studies [140, 155] found benefit of surgery in terms of reduced mortality but no improvement in neurological outcome.

Conservative treatment is suggested either for the least severe patients or for those considered to be beyond possible treatment [156]. Some authors suggested that, in patients with acute subdural hematoma for whom the neurosurgeon sees no clear advantage of acute surgery over conservative management, an initial conservative approach may be considered [139]. Hematoma size and mass effect were associated with failure of conservative management in some trials [157,158] but not in others [159].

Recommendations

- **R 7.1: The experts suggest that the emergency surgical evacuation of acute subdural hematoma can improve the neurological outcome in TBI patients under the following conditions:**
 - o age <65 years OR 65–80 years but with low frailty score

AND

- o impaired consciousness (GCS ≤ 8 AND/OR loss of ≥ 2 GCS points) not explained by another mechanism OR intracranial hypertension refractory to medical treatment;

AND

- o ASDH with thickness >10 mm AND/OR midline shift >5 mm

EXPERT OPINION (STRONG AGREEMENT)

- **R 7.2: The experts suggest that later or less invasive surgery should be considered after ASDH if rebleeding is documented in chronic SDH.**

EXPERT OPINION (STRONG AGREEMENT)

- **R 7.3: The experts suggest that conservative treatment should be considered in patients with ASDH outside the above circumstances. Follow-up with brain CT is recommended:**
 - o urgently if clinical course deteriorates;
 - o at an early stage (7–10 days) if antiplatelet or anticoagulant treatment needs to be resumed;
 - o at a later stage (3–4 weeks) in other cases.

EXPERT OPINION (STRONG AGREEMENT)**4. DEPRESSED SKULL FRACTURE AND SKULL BASE FRACTURE ASSOCIATED WITH CEREBROSPINAL FLUID LEAKAGE**

Edouard SAMARUT, Clémentine GALLET, Alexandre BANI-SADR, Marion LE MARECHAL, Romain MANET

Question 8: In TBI patients with depressed skull fracture, does surgical management improve neurological outcome?

In the absence of intracranial lesions, depressed skull fracture has good prognosis [160–164]. Dural tears must be excluded in case of open depressed skull fracture, which otherwise corresponds to penetrating TBI (pTBI) (see section 5). In a retrospective series of 128 patients treated for TBI associated with depressed skull fracture, the depth of the depressed skull fracture (OR 1.3; 95% CI 1.14–1.46), presence of pneumocephalus (OR 2.5; 95% CI 1.05–5.76) and adjacent contusion or hematoma (OR 5.5; 95% CI 1.57–8.82) were predictive of dural tear [165].

According to the Brain Trauma Foundation guidelines [166], neurosurgical indications include open depressed skull fracture associated with a contaminated wound, signs of local infection, uncontrollable bleeding, intracranial contusion or hematoma, cosmetic damage, signs of dural effraction, mass effect on the cerebral parenchyma, or depression depth >1 cm below the internal table. However, two studies, including 67 patients [162] and 232 patients [167], found no significant improvement in outcome in terms of neurological results, epilepsy or infection in patients treated surgically according to the BTF criteria compared with patients treated conservatively.

For uncontaminated skull fracture, wound excision and closure can be performed, not necessarily by a neurosurgeon. If neurosurgery is indicated, titanium osteosynthesis is preferred for reconstruction [168–170]. Depressed skull fracture is not an absolute emergency. Surgical treatment within 24 h is recommended [171].

Depressed skull fracture in the frontal sinus is associated with cosmetic damage, CSF rhinorrhea, short-term infectious complications (meningitis, empyema) and long-term complications (mucocele). In practice, two parameters need to be considered: integrity of the dura in the posterior wall, and integrity of the nasofrontal ducts. The frontal sinus should be preserved in the absence of major posterior wall defect, obvious dural tear or damage to the nasofrontal ducts. Otherwise, the procedure should be completed by obliteration or cranialization of the frontal sinus [171–175].

In depressed skull fracture exerting a mass effect on a venous sinus, several case reports [160,176–179] and a series of 13 cases [180] reported surgical treatment (direct fracture reduction, decompressive craniectomy, or CSF drainage) in cases of refractory intracranial hypertension.

Depressed skull fracture increases the risk of post-traumatic epilepsy (PTE) (OR 1.88, 95% CI 0.92–3.80; $p = 0.081$) [181]. PTE was reported in 16% of 333 patients in a retrospective cohort of patients treated for closed depressed skull fracture [182], and was related to initial clinical severity [162,181,182]. Although PTE was an independent factor for poor outcome [181], it was not influenced by surgical modalities for depressed skull fracture or antiepileptic prophylaxis [167,181]. Antiepileptic prophylaxis is therefore rarely prescribed in Europe ($<20\%$ of responders), in contrast to other parts of the world [183].

Recommendations

- **R 8.1: In a patient with traumatic brain injury associated with a depressed skull fracture, the experts suggest prompt neurosurgical management, within 24 h, in any of the following circumstances:**
 - complex or contaminated wound or evidence of local infection;
 - suspected dural tear/pneumocephalus;
 - CSF leakage;
 - significant mass effect on brain parenchyma;
 - other neurosurgical lesion(s);
 - major cosmetic damage.

EXPERT OPINION (STRONG AGREEMENT)

- **R 8.2: In a patient with traumatic brain injury associated with a depressed skull fracture involving the frontal sinus, the experts suggest surgical management with anterior wall reduction/osteosynthesis alone in the following circumstances:**
 - large posterior wall defect;
 - obvious dural tear;
 - nasofrontal canal involvement.
- Otherwise, the procedure must be completed with cranialization of the frontal sinuses.

EXPERT OPINION (STRONG AGREEMENT)

- **R 8.3: In a patient with traumatic brain injury associated with depressed skull fracture and epileptic seizure, experts suggest initiating secondary antiepileptic prophylaxis. Primary antiepileptic prophylaxis should not be systematically prescribed in non-penetrating TBI associated with depressed skull fracture.**

EXPERT OPINION (STRONG AGREEMENT)

Question 9. In a patient with traumatic brain injury associated with skull base fracture and cerebrospinal fluid leakage, does surgical management improve neurological outcome?

Post-traumatic skull base cerebrospinal fluid (CSF) leakage occurs in 0.3%–39% of skull base fractures [184–186]. More than 50% of traumatic leaks occur within the first 48 h and 70% within the first 7 days. In the case of "true" tears, they occur almost systematically after 3 months [184,187–189]. Leakage mainly involves the frontal sinus (30% of cases), the sphenoidal sinus (11–30 %) or the ethmoid sinus (15–19%) [190,191]. Meningitis occurs in 0.2% of cases within the first 24 h after trauma, in 9% after the first week, and in 19% after the second week. The cumulative risk of meningitis is estimated to be about 1% per day for the first 2 weeks, then 7% per week for the first month [192,193]. Neurological outcome is mainly determined by the initial severity of TBI and not by the presence of CSF leakage [193,194].

Surgery is the preferred option if the leak persists after 3–7 days' well-conducted medical management [193–196]. Surgery reduces the overall risk of meningitis from 30% before surgery to 4% after. The 10-year cumulative risk also decreases, from 85% before surgery to 7% after. Recurrence despite surgery is estimated to be between 12.5% and 17%, with 1.3% mortality [197,198].

In case of associated intracranial lesions, craniotomy is preferred to endoscopic endonasal repair of the dural defect [185,193,196,199–201]. In cases where emergency intracranial surgery is required for the initial trauma, some authors suggest closing the dural defect in the same step in the absence of intracranial hypertension (ICH). In case of ICH, a two-stage approach is recommended [193,194,201].

Small dural defects (diameter ≤ 1 cm) can be treated with an endonasal approach, but larger defects (>1 cm) require craniotomy [185,193,194,196,201–203]. The location of the defect is an additional factor in the choice between an endonasal approach or craniotomy: medial defects can be easily treated with an endonasal approach, whereas lateral, very anterior or intermediate defects require craniotomy [185,193,194,196,199–202,204].

Conservative treatment for post-traumatic skull-base CSF leakage includes bed rest (bedhead at 30 °) and laxatives, antitussives and antiemetics for at least 48–72 h. CSF leakage can thereby be stopped in 4–5 days in 60% of cases, and in 7 days in 85% [187,193,196,198,205]. In case of failure, 7–10 days' external (lumbar or ventricular) CSF drainage can be considered, with a target flow rate of 10–15 mL/hour. In most patients, this procedure was successful [195,205,206]. Post-operative external lumbar drainage can reduce the duration of rhinorrhea [207,208] although there is no evidence of impact on neurological outcome or risk of infection.

The SFAR/SPILF 2023 guidelines [209] do not recommend antibiotic prophylaxis for skull-base fractures. Antibiotic prophylaxis should be given for neurosurgical procedures only.

No studies compared the risk of long-term meningitis after dural tear in patients having received vaccination versus no vaccination. The epidemiology of meningitis cases was 50% *Streptococcus pneumoniae* and 20% *Haemophilus influenzae* [210].

The radiological location of a post-traumatic skull-base dural tear can be challenging. MRI and combined CT/MRI are more sensitive than conventional CT [211]. In a recent systematic review of the detection of CSF rhinorrhea [212], the sensitivity and specificity of the different detection methods were as follows: high-resolution CT: 0.93/0.50; MRI: 0.94/0.77, CT myelography: 0.95/1.00; isotopic cisternography: 0.90/0.50; MRI cisternography with contrast enhancement: 0.99/1.00; nasal endoscopy: 0.58/1.00; intrathecal fluorescein: 0.96/1.00.

Recommendations

- **R 9.1 In a patient with traumatic brain injury associated with skull-base fracture and cerebrospinal fluid leakage, the experts suggest prompt surgery (within 24 h) in case of major dural defect associated with abundant liquorrhea, to improve the neurological prognosis. If neurosurgery is indicated for other associated lesions, closure of the dural defect in the same step is not recommended in case of intracranial hypertension.**

EXPERT OPINION (STRONG AGREEMENT)

- **R 9.2: In a patient with traumatic brain injury associated with skull-base fracture and cerebrospinal fluid leakage, the experts suggest surgery in case of non-abundant liquorrhea refractory to conservative treatment for more than 7 days.**

EXPERT OPINION (STRONG AGREEMENT)

- **R 9.3: In a patient with traumatic brain injury associated with skull-base fracture and cerebrospinal fluid leakage, the experts suggest that conservative treatment of post-traumatic dural tears of the skull base should be combined with bed-rest at 30 ° and laxatives, cough suppressants and antiemetics for at least 72 h.**

EXPERT OPINION (STRONG AGREEMENT)

- **R 9.4: In a patient with traumatic brain injury associated with skull-base fracture and cerebrospinal fluid leakage, the experts recommend that vaccinations against pneumococcus, Haemophilus influenzae and meningococcus should be carried out or updated, to reduce the risk of meningitis:**
 - pneumococcal vaccination: 15-valent conjugate vaccine for under-18 year-olds (1 dose for over-2 year-olds) or 20-valent vaccine for over-18 year-olds (1 dose not followed by a non-conjugate vaccine);
 - Haemophilus influenzae vaccination (1 dose);
 - Meningococcus B and ACYW vaccination: catch-up according to general recommendations (meningococcal ABCYW before 2 years of age and meningococcal ACYW before 25 years of age).

EXPERT OPINION (STRONG AGREEMENT)

- **R 9.5:** In a patient with traumatic brain injury associated with skull-base fracture and cerebrospinal fluid leakage, the experts suggest screening for traumatic dural defect on bone-window CT scan with millimeter slices and MRI with high-resolution 3D T2 sequences. If in doubt, a CT-myelography can be performed, followed, if negative, by isotope cisternography, to improve diagnostic performance.

EXPERT OPINION (STRONG AGREEMENT)

5. PENETRATING TRAUMATIC BRAIN INJURY

Arnaud DAGAIN, Christophe JOUBERT, Sébastien GAZZOLA, Pierre ESNAULT, Marion LE MARECHAL

Question 10. In a patient with penetrating traumatic brain injury, under what circumstances is neurosurgical treatment likely to improve neurological prognosis?

Penetrating traumatic brain injury (TBI) is associated with high mortality, up to 85% [213–219]. Aggressive management of these patients improves survival [218,220–222].

The following factors have been reported to be associated with high mortality: age over 35 years, unresponsive pupils, respiratory arrest or hypotension on admission, and bi-hemispheric (excluding bi-frontal), trans-hemispheric, transventricular or posterior fossa ballistic trajectories [214,219,223,224]. Factors moderately associated with higher mortality included ICP > 20 mm Hg and Glasgow Coma Score (GCS) 3 or 4 at presentation [223]. On multivariate analysis, significant factors for mortality were bullet trajectory and pupillary response [223].

The 2 main predictive scores for mortality are the Maritzburg score and the SPIN score [225,226]. The Maritzburg score is based on 2 criteria: GCS < 5 and externalization of the brain through the skin wound; presence of both has a positive predictive value of 92% (sensitivity 73%, specificity 98%). The SPIN score is based on a motor GCS subscore (mGCS), pupillary reactivity, absence of self-inflicted injury, transfer to a referral center, female gender, lower Injury Severity Score (ISS) and lower International Normalized Ratio (INR) (see Appendices), all shown to correlate strongly with survival [226]. These scores can be used for early prognostication, but should not be used in isolation to contraindicate aggressive management.

GCS at initial treatment is the most important clinical prognostic factor [213]. In particular, a GCS < 5 is generally associated with very poor prognosis. Patients operated on for penetrating TBI with initial GCS 6–8 had better neurological prognosis than those treated conservatively (24% vs 0%) [227]. The rate of good neurological outcome in patients operated on with GCS ≥ 9 was high, at 77.8% [228]. These data suggest that neurosurgery should be widely used for patients with penetrating TBI.

Ideally, surgery should be performed within 5 h of trauma to reduce the risk of cerebro-meningeal infection [229,230,231]. In the absence of intracranial hypertension, surgery consists in plane-by-plane debridement, including watertight dural closure and skin closure if the scalp is not devitalized, to reduce the risk of infection, mainly from CSF fistulation [231–235]. Removal of bone fragments or bullets at a distance from the entry point, particularly in eloquent areas of the brain, is not systematic: it may reduce the risk of post-traumatic epilepsy [236], but at the cost of increased functional morbidity [231,232,235,237].

In case of extensive lesions or intracranial mass effect, surgery, combined with control of intracranial pressure and systemic aggression by intensive care, is part of a 'neuro-damage control' strategy aimed at reducing secondary cerebral lesions [238]: in addition to plane-by-plane debridement, evacuation of compressive hematomas and removal of accessible foreign bodies [229], or even decompressive craniectomy [239,240] with watertight dural closure, ideally with autologous augmentation plasty (galea or facia lata) if necessary [241,242].

Neurological outcome after superficial debridement was very heterogeneous, and essentially correlated with initial neurological severity

as reflected by GCS [228,243]: the rate of good neurological prognosis varied from 0.8% to 75% [213,215,223,227], with 0% for an initial GCS 6–8, 50% for 9–11, and 95% for ≥ 12 [227]. Superficial debridement was therefore appropriate in selected patients with GCS ≥ 12, but offered limited benefit in more severe cases with intracranial hypertension, indicating more aggressive debridement in the latter.

Incidence of epileptic seizures in penetrating TBI within the first 7 days was high, ranging from 14% to 20% [244]. Patients with penetrating TBI were 3 times more likely to be readmitted to hospital within 2 years for concomitant seizure than patients with closed TBI [245,246]. Although incidence of early seizures was higher with penetrating TBI, incidence of late seizures did not differ [247]. Finally, incidence of epilepsy was unchanged by initial surgical treatment [242]. There are no randomized controlled trials of anticonvulsant prophylaxis in patients admitted for penetrating TBI; it appears to be acceptable in the first week, but not thereafter [247,248].

The SFAR / SPILF 2023 expert recommendations advocated antibiotic prophylaxis with 2 g amoxicillin-clavulanic acid, which can be continued for 24–48 h in penetrating TBI with a soiled wound [209].

No studies compared the long-term risk of meningitis with or without vaccination. The epidemiology of meningitis in cases of dura mater breach was 50% *Streptococcus pneumoniae* and 20% *Haemophilus influenzae* [210].

Recommendations

- **R 10.1:** In a patient presenting with penetrating brain injury, experts suggest that the risk of mortality (SPIN score, Maritzburg score) should be assessed and taken into account, in a non-isolated manner, in the indication for emergency surgery.

EXPERT OPINION (STRONG AGREEMENT)

- **R 10.2:** In a patient presenting with penetrating brain injury, associated with Glasgow Coma Score ≥ 5, the experts suggest urgent neurosurgical management.

EXPERT OPINION (STRONG AGREEMENT)

- **R 10.3:** In a patient presenting with penetrating brain injury, associated with a Glasgow coma score ≤ 4, in absence of bilateral non-reactive mydriasis and of pejorative radiological criteria, the experts suggest evaluating the need for urgent neurosurgical management individually.

EXPERT OPINION (STRONG AGREEMENT)

- **R 10.4:** In a patient presenting with penetrating brain injury, the experts suggest that, in case of significant cortical damage, primary antiepileptic prophylaxis should be given for at least 7 days.

EXPERT OPINION (STRONG AGREEMENT)

- **R 10.5:** In a patient presenting with penetrating brain injury, the experts suggest that vaccinations against pneumococcus, meningococcus, *Haemophilus influenzae* and tetanus should be carried out or updated:
 - o in people who are not up-to-date: anti-tetanus vaccination combined with injection of 250 IU of anti-tetanus human immunoglobulin in case of extensive penetrating wound with foreign body or wound treated late;
 - o pneumococcal vaccination: 15-valent conjugate vaccine for under-18 year-olds (1 dose for over-2 year-olds) or 20-valent vaccine for over-18 year-olds (1 dose not followed by non-conjugate vaccine);

- o Meningococcus B and ACYW vaccination: catch-up according to general recommendations (meningococcus ABCYW before age 2 and meningococcus ACYW before age 25);
- o Haemophilus influenzae vaccination (1 dose)

EXPERT OPINION (STRONG AGREEMENT)

Question 11. In a patient presenting with penetrating brain injury, when and how should screening for intracranial vascular lesions be performed?

Traumatic intracranial vascular lesions have been identified in up to 60% of patients with penetrating TBI [249,250].

Post-traumatic intracranial aneurysm (PTIA) is common in penetrating TBI, with incidence between 20% and 50% [214,218]. It is difficult to diagnose, associated with high mortality, and also poses the problem of choosing a treatment modality.

PTAI may occur in the first hours after trauma or later (several days to several weeks). Progression is uncertain, but PTAI is associated with intracranial hemorrhage in 80% of cases and subdural hematoma in 26% [214]. Ruptured PTAI is associated with high mortality, up to 50% [219].

Given the significant morbidity and mortality associated with failure to detect and treat PTAI, several authors recommended systematic screening for penetrating TBI [213,218,241]. Cerebral arteriography is the gold-standard for early detection of these lesions in penetrating TBI; cerebral CT-angiography has lower sensitivity [251,252]. Criteria for performing digital subtraction arteriography were defined by Bell [213,218,241]:

- o penetrating injury through the peritonsillar and orbitofrontal regions;
- o known cerebrovascular lesion with or without pseudoaneurysm observed on initial exploration;
- o blast injury with Glasgow score <8 (closed or penetrating);
- o evidence of vasospasm on transcranial Doppler;
- o Spontaneous unexplained decrease in cerebral tissue oxygen pressure (PtiO₂).

To these criteria, the following may be added:

- o penetrating trajectory close to the main arterial axes;
- o multiple dural injuries;
- o intracerebral hematoma.

Early vascular imaging may be negative either because the lesion is too small to be detected on CT-angiography or because it appears to be secondary. If in doubt, follow-up imaging in 2–3 weeks should be considered.

Definitive multimodal treatment should be started immediately [214,219]. There is no evidence that surgical treatment is superior to endovascular treatment [241,253–255].

Recommendations

- **R 11.1: In a patient with a penetrating brain injury, the experts suggest systematic screening for intracranial vascular lesions on CT-angiography.**

EXPERT OPINION (STRONG AGREEMENT)

- **R 11.2: In a patient presenting with penetrating brain injury, the experts suggest that this examination be supplemented with a 6-axis digital subtraction cerebral arteriogram in case of one or more of the following:**
 - o detection or suspicion of a vascular lesion on CT-angiography;

- o penetrating injury in the periorbital and/or fronto-orbital region;
- o multiple dural injuries;
- o trajectory and/or hematoma close to the major vascular axes;
- o penetrating or blast TBI with Glasgow score <8;
- o detection of vasospasm on transcranial Doppler and/or spontaneous unexplained decrease in cerebral tissue oxygen pressure (PbtO₂).

EXPERT OPINION (STRONG AGREEMENT)

- **R 11.3: In patients presenting with penetrating TBI, associated with symptomatic or asymptomatic intracranial vascular injury at high risk of rupture and/or neurological worsening, the experts suggest that urgent treatment should be discussed.**

EXPERT OPINION (STRONG AGREEMENT)

- **R 11.4: No recommendations on modality (endovascular or surgical), which should be discussed collegially.**

NO RECOMMENDATION

- **R 11.5: In patients presenting with penetrating TBI, the experts suggest that, in the absence of proof of an initial intracranial vascular lesion, cerebrovascular imaging should be repeated approximately 14 days after trauma.**

EXPERT OPINION (STRONG AGREEMENT)

6. POST-TRAUMATIC CEREBROSPINAL FLUID DISORDERS

Romain MANET, Stéphane GOUTAGNY, Baptiste BALANCA, Vincent JECKO, Philippe DECQ, Arnaud DAGAIN

Question 12: In a patient with traumatic brain injury associated with post-traumatic peri-encephalic collections of cerebrospinal fluid, under what circumstances is surgical intervention likely to improve neurological outcome?

The pathophysiology of extra-axial cerebrospinal fluid (CSF) collections in adults is poorly understood. In practice, two entities can be distinguished. On the one hand, hygroma is thought to result from tearing of arachnoid layer adhesions to the dura mater [256,257] or, more rarely, from traumatic or iatrogenic rupture of an arachnoid cyst [258–261], leading to sequestration of CSF in the resulting subdural space. This passive accumulation, favored by post-traumatic atrophy [262], does not interfere with normal CSF circulation. Intracranial pressure (ICP) and neurological examination generally remain normal. The vast majority of hygromas are benign and are treated conservatively [263]. Conversely, external hydrocephalus involves 'active' sequestration of CSF within the subarachnoid spaces (SAS) as a result of resorption failure, favored by post-traumatic subarachnoid hemorrhage [264,265], which generally leads to an increase in ICP [266] and worsens the neurological condition of the patient [264]. Extra-axial CSF collection after decompressive craniectomy is very common and is often interpreted as hygroma [267–270], but may be more consistent with external hydrocephalus, with a risk of chronic post-traumatic hydrocephalus requiring definitive CSF shunting [264,267,268,271].

Management of peri-encephalic post-traumatic CSF collection has not been the focus of any guidelines. At the 2019 Seattle International Consensus Conference on the Management of Severe TBI, external lumbar drainage of CSF was not selected as a therapeutic measure for the management of post-traumatic intracranial hypertension [11]. Our systematic review of the literature on external lumbar drainage in TBI retrieved 10 original articles [265,272–280]. Three studies with interim results included in later publications [281–283] and 1 published in Hungarian [284] were not included in our analysis. Overall, our analysis included 221 patients with TBI, the majority of whom were adults. In 3 studies, no patients had previously received external ventricular

drainage [275,277,278], which was performed systematically in 4 studies [272,276,279,280]; in the other 3 studies, it was used optionally. The weighted mean absolute and percentage reductions in ICP following external lumbar drainage were -17 mmHg and -59% , respectively. The meningitis rate was 6% ($12/208$), with tonsillar involvement in 7% of patients ($21/295$), primarily in studies where drainage protocols included counter-pressures of 0 or $+5$ cmH₂O (relative to the foramen of Monro), without the possibility of assessing the associated morbidity and mortality. To mitigate this risk, all authors relied on CT criteria, including the patency of the basal cisterns (particularly the prepontine and/or quadrigeminal cisterns), absence of lesions with significant mass effect, absence of midline shift >5 mm or >10 mm, absence of uncus or tonsillar herniation, and the 'Innsbruck intracranial reserve score,' which incorporates several of these elements [275]. In addition, one study proposed including a criterion of external hydrocephalus, defined by clinical features (delayed onset of intracranial hypertension and poor response to osmotherapy) and CT findings (progressive and paradoxical dilatation of the subarachnoid spaces, including the sylvian fissures, interhemispheric fissures and cortical sulci) [265]. Finally, to reduce the risk of a pressure gradient between the cranial and spinal compartments, the zero reference level for external lumbar drainage was set at the foramen of Monro (or the external auditory canal) in all studies.

Recommendations

Experts suggest distinguishing 2 types of post-traumatic periencephalic cerebrospinal fluid (CSF) collection:

- o generally stable (rarely progressive) unilateral collection AND absence of neurological signs AND normal intracranial pressure = hygroma;
- o progressive collection on successive CT scans, generally bilateral AND neurological worsening AND/OR increased intracranial pressure = external hydrocephalus
- R 12.1: In patients presenting TBI associated with hygroma, the experts suggest conservative treatment in first line.

EXPERT OPINION (STRONG AGREEMENT)

- R 12.2: In patients presenting TBI associated with external hydrocephalus, experts suggest cerebrospinal fluid drainage, either by lumbar puncture or by external lumbar drainage, after CT confirmation of:
 - o patency of the basal cisterns,
 - o AND absence of midline deviation >10 mm,
 - o AND absence of tonsillar involvement.

In case of intracranial hypertension, this option should only be considered after 1st-line measures have failed.

EXPERT OPINION (STRONG AGREEMENT)

- R 12.3: In patients presenting traumatic brain injury associated with external hydrocephalus treated by external lumbar drainage, the experts suggest placing the external lumbar drainage zero reference at the external auditory canal. In addition, in case of intracranial hypertension, intracranial pressure should be continuously monitored and the reference counter-pressure should not be lowered below 10 mmHg. Finally, if the pressure gradient between intracranial pressure and lumbar CSF pressure is greater than 5 mmHg, lumbar drainage should be discontinued.

EXPERT OPINION (STRONG AGREEMENT)

7. SPECIFICS OF NEUROSURGICAL MANAGEMENT OF PEDIATRIC TRAUMATIC BRAIN INJURY

Nathalie CHIVORET, Andres COCA, Guillaume MORTAMET, Matthieu VINCHON

Question 13. In neonates less than 1 month of age with

traumatic brain injury associated with extradural hematoma, under what circumstances is surgical intervention likely to improve neurological outcomes?

Neonatal extradural hematoma (EDH) is a recognized complication of forceps and vacuum-assisted delivery, but is rare. In recent studies, incidence of intracranial hemorrhage following instrumental delivery was 0.4 – 1% , and the incidence of neonatal EDH can be presumed to be considerably lower [285].

Neurological status at presentation, assessed on the Child Coma Scale (CCS) or GCS, is strongly prognostic. Children with CCS or GCS scores below 8 are subject to high mortality. In reported series, EDH was predominantly located in the parietal and temporoparietal regions, but no correlation was found between location and prognosis [286,287]. According to literature reviews, skull fracture does not significantly impact prognosis, including GOS score [287,288]. Neonates and infants are more resilient to acute elevations in ICP, owing to anatomical characteristics such as unfused cranial suture, open fontanelle, and relatively larger extracerebral spaces. Moreover, in contrast to adults, pediatric EDH is more frequently of venous origin [289].

It is generally accepted that many EDHs can be managed conservatively without need for surgery. This is especially true for neonatal EDH, which often resolves spontaneously. In neonates, EDH tends to liquefy more rapidly than in older age groups, and most analyses of contents found predominantly liquid blood [290,291].

A conservative approach is typically favored in patients with small hematoma (<20 mm in thickness), in the absence of midline shift or clinical deterioration, and with good neurological status (high GCS). Such cases require close monitoring in an intensive care setting, including regular neurological examinations and repeat imaging if deterioration is suspected.

The main indications for surgical management of EDH in neonates comprise substantial hematoma thickness, significant brain displacement, and depressed skull fracture or associated hydrocephalus. A review of the literature suggested that hematoma volume >30 mL, thickness >15 mm and midline shift >5 mm are strong indicators for surgical intervention [121,292]. The standard treatment is surgical evacuation, typically via craniotomy. However, in neonates, craniotomy is particularly invasive and associated with significant risks. Whenever feasible, less invasive approaches are preferred, to minimize surgical morbidity and optimize hematoma management. In cases where EDH is associated with a cephalohematoma, aspiration of the cephalohematoma may be considered. Alternatives such as skull fracture aspiration or direct epidural aspiration may be appropriate, particularly in EDH without associated skull fracture or cephalohematoma [293–295]. Craniotomy should be reserved for cases where less invasive techniques fail. In cases of liquefied hematoma, evacuation through a single burr-hole may be sufficient. If craniotomy is performed, osteosynthesis of the bone flap should be avoided, due to potential interference with subsequent skull growth.

Ongoing clinical and neurological monitoring is essential for managing EDH in infants under 1 year of age. These patients require close observation in an intensive care setting, with repeated neurological assessments and follow-up imaging when clinically indicated, mindful of the risks associated with cumulative ionizing radiation from CT scans.

Recommendations

- R 13.1: In neonates with traumatic brain injury associated with extradural hematoma, the experts suggest emergency surgical evacuation in case of clinical or radiological signs of severity (>5 mm deviation of the midline or brainstem compression).

EXPERT OPINION (STRONG AGREEMENT)

- R 13.2: As there is no consensus on the optimal surgical technique, the experts suggest that less invasive approaches be considered before resorting to craniotomy. These strategies

include aspiration of the cephalohematoma or other methods, such as puncture through a skull fracture or epidural puncture, particularly in cases of extradural hematoma without associated skull fracture or cephalohematoma.

EXPERT OPINION (STRONG AGREEMENT)

- **R 13.3:** In neonates with traumatic brain injury associated with extradural hematoma, experts recommend a conservative approach in the absence of clinical or radiological signs of severity. In these cases, close monitoring in an intensive care unit is essential, including regular clinical examination and follow-up imaging.

EXPERT OPINION (STRONG AGREEMENT)

Question 14. In infants under 2 years of age with non-accidental traumatic brain injury associated with subdural hematoma, under what circumstances is surgical treatment likely to improve neurological prognosis?

Non-accidental infantile TBI, commonly known as "shaken baby syndrome," is often associated with subdural hematoma (SDH), particularly in infants. Infantile SDH presents as a mixture of blood and cerebrospinal fluid; it is a frequent condition at this age and has no direct equivalent in adult pathology. Although often mistakenly referred to as chronic SDH, it typically develops within a few days of trauma and follows a relatively acute course. Prognosis correlates primarily with initial clinical severity ($p = 0.001$) and is not significantly influenced by shunt-related complications ($p = 0.27$) [296,297]. Overall incidence of complications was significantly associated with initial clinical severity ($p = 0.013$), preoperative hospital stay time ($p = 0.016$), and time between trauma and surgery ($p = 0.037$). Several studies confirmed that infant maltreatment and initial clinical severity, as indicated by stay in intensive care, are important independent prognostic factors [298]. Moreover, the morbidity and mortality associated with non-accidental pediatric TBI (maltreatment) are higher than in accidental TBI [298].

The primary goal of treatment is to relieve intracranial hypertension and prevent complications associated with SDH, such as cerebral atrophy, visual disorder and craniocerebral disproportion. Neurosurgical indications for emergency evacuation of the hematoma are typically based on clinical criteria, including GCS score ≤ 12 at admission, bulging fontanel or other signs of intracranial hypertension (e.g., on transcranial Doppler), which may improve neurological prognosis [297–300]. Initial treatment for intracranial hypertension may involve transfontanelar subdural puncture. However, intracranial hypertension is often only temporarily controlled by subdural puncture, which may need to be repeated. Although there is no definitive consensus on surgical techniques specific to non-accidental trauma, in retrospective studies analyzing large cohorts, subdural-peritoneal shunting, in addition to subdural puncture, was the most commonly employed treatment for SDH [296,297]. The role of decompressive craniectomy has been widely studied in accidental TBI [301–303], but results in infants with non-accidental TBI are uncertain, precluding any recommendation. The therapeutic window for intervention is especially limited in toddlers, and the presence of a 'big black brain' at admission is a contraindication. Other patients can typically be managed with intensive resuscitation, early monitoring of ICP, and CSF withdrawal via lumbar puncture if the cisterns are patent, or subdural drainage if they are not. Conservative treatment may be considered in cases of small (<10 mm), clinically well-tolerated SDH (including isolated seizure), with no signs of intracranial hypertension. However, there are no direct medical treatments for SDH [297]. Acute SDH, consisting of fresh clotted blood, comparable to the lesion seen in adults, is very rare in infants. The literature on this subject is extremely limited.

Recommendations

- **R 14.1:** In infants with non-accidental traumatic brain injury associated with subdural hematoma, the experts suggest rapid surgical management, preferably by drainage.

EXPERT OPINION (STRONG AGREEMENT)

- **R 14.2:** No recommendations are made on decompressive craniectomy in non-accidental traumatic brain injury in infants.

NO RECOMMENDATION

- **R 14.3:** In infants with non-accidental traumatic brain injury associated with subdural hematoma, the experts suggest considering conservative management for chronic subdural hematomas with small thickness (<10 mm) that are clinically well-tolerated, accompanied by close surveillance and regular imaging monitoring.

EXPERT OPINION (STRONG AGREEMENT)

Question 15. In infants under 2 years of age with ping-pong fracture, under what circumstances is surgical management likely to improve neurological outcome?

Delivery-related injuries include ping-pong fracture, occurring in 1–2.5 cases per 10,000 live births. This is characterized by a bony depression in the cranial vault without loss of continuity, and is rarely associated with intracranial lesions. There are no clearly identified predictive factors to determine which cases will resolve spontaneously and which will require intervention.

A recent systematic review, including 54 articles and 228 children under 20 months of age (excluding complex fractures associated with intracranial lesions) [304], reported that 30% of children received surgical management, 30% underwent interventional management (aspiration), and 40% were managed conservatively. The review found no significant difference in neurological outcome according to type of management, with favorable prognosis (no neurological deficit or epilepsy) in more than 96% of patients. The surgical indications reported were as follows:

- o presence of a foreign body;
- o wound contamination, infection or need for trimming;
- o neurological deficit;
- o signs of intracranial hypertension;
- o unsuccessful correction by external maneuver (aspiration);
- o risk of difficulty with close neurosurgical follow-up of conservative treatment;
- o intracranial hematoma;
- o hygroma;
- o major cosmetic concerns.

The authors recommended that infants with symptomatic ping-pong fracture undergo ultrasound examination, and those with an intracranial lesion visible on ultrasound or depression >1 cm undergo CT. Several of the included studies [304], as well as a recent retrospective series of 7 cases [305], advocated conservative treatment for fractures through external maneuvers (e.g., suction).

While the possibility of fracture reduction is emphasized, there is insufficient data on the impact of these procedures on neurological outcome. Consequently, the effectiveness of the maneuver remains uncertain.

Recommendations

- **R 15.1:** In infants presenting ping-pong fracture, the experts suggest emergency surgery in the following situations:
 - o intracranial hypertension;
 - o significant mass effect on the parenchyma or intracranial hematoma or peri-encephalic CSF collection.

EXPERT OPINION (STRONG AGREEMENT)

- **R 15.2: In infants presenting ping-pong fracture, the experts suggest that, in the absence of the above criteria, management should initially be conservative. Surgical management may be re-evaluated in the absence of favorable outcome.**

EXPERT OPINION (STRONG AGREEMENT)

Coordinators

Romain MANET (SFNC), Arnaud DAGAIN (SFNC)

Methodological supervision

Jean François PAYEN (SFAR, ANARLF), Hugues DE COURSON (CRC SFAR, ANARLF)

Experts

French Society of Neurosurgery (SFNC): Cyrille CAPEL (Amiens), Arnaud DAGAIN (Toulon), Philippe DECQ (Paris), Matthieu FAILLOT (Paris), Clémentine GALLET (Lyon), Vincent JECKO (Bordeaux), Stéphane GOUTAGNY (Paris), Christophe JOUBERT (Toulon), Romain MANET (Lyon), Édouard SAMARUT (Nantes)

French Society of Pediatric Neurosurgery (SFNCP): Nathalie CHIVORET (Nice), Andres COCA (Strasbourg), Mathieu VINCHON (Lyon)

French Society of Private-Practice Neurosurgeons (SFNCL): Alice ROLLAND (Montpellier)

French Society of Anesthesia, Critical Care and Perioperative Medicine (SFAR), and Neurocritical Care and French-Speaking Neuro Anesthesiology Society (ANARLF): Baptiste BALANCA (Lyon), Mickael CARDINALE (Toulon), Hugues DE COURSON (Bordeaux), Pierre ESNAULT (Toulon), Jean-Denis MOYER (Caen), Jean François PAYEN (Grenoble), Hervé QUINTARD (Genève), Stéphanie SIGAUT (Paris)

French-Speaking Group of Emergency and Intensive Care in Pediatrics (GFRUP): Guillaume MORTAMET (Grenoble)

French Society of Neuroradiology (SFNR): François COTTON (Lyon), Alexandre BANI-SADR (Lyon), Sébastien GAZZOLA (Toulon)

French-speaking Society of Infectious Diseases (SPILF): Marion LE MARECHAL (Grenoble)

French Society of Physical Medicine and Rehabilitation (SOFMER): Jacques LUAUTÉ (Lyon), Eric VERIN (Rouen)

Work groups

Prognostic factors to be considered when indicating emergency craniotomy in adult patients with traumatic brain injury

Romain MANET, Mickaël CARDINALE, François COTTON, Hervé

QUINTARD, Jacques LUAUTE, Éric VERRIN, Arnaud DAGAIN

Extradural hematoma

Cyrille CAPEL, Jean Denis MOYER

Acute subdural hematoma

Matthieu FAILLOT, Stéphanie SIGAUT, Alice ROLLAND

Depressed fractures and skull base fractures associated with cerebrospinal fluid fistula

Edouard SAMARUT, Clémentine GALLET, Alexandre BANI-SADR, Marion LE MARECHAL, Romain MANET

Penetrating traumatic brain injury

Arnaud DAGAIN, Christophe JOUBERT, Sébastien GAZZOLA, Pierre ESNAULT, Marion LE MARECHAL

Post-traumatic cerebrospinal fluid disorders

Romain MANET, Stéphane GOUTAGNY, Baptiste BALANCA, Vincent JECKO, Philippe DECQ, Arnaud DAGAIN

Specifics of neurosurgical management of pediatric traumatic brain injury

Nathalie CHIVORET, Andres COCA, Guillaume MORTAMET, Matthieu VINCHON

Reading groups

SFNC board

Claire Haegelen (President); Ilyess Zemmoura (Vice-president); Damien Bresson (Secretary); Olivier Hamel (Deputy secretary); Thomas Roujeau (Treasurer); Hayat Belaid; Marc Baroncini

SFAR board

Jean-Michel Constantin (President); Marc Léone (1st vice-president); Karine Nouette-Gaulain (2nd vice-president); Isabelle Constant (General secretary); Frédéric Le Saché (Deputy general secretary); Evelyne Combettes (Treasurer); Olivier Joannes-Boyau (Deputy treasurer); Pierre Albaladejo; Julien Amour; Hélène Beloeil; Valérie Billard; Marie-Pierre Bonnet; Julien Cabaton; Sébastien Campard; Vincent Collange; Marion Costecalde; Violaine D'Ans; Laurent Delaunay; Delphine Garrigue; Frédéric Lacroix; Sigismund Lasocki; Anne-Claire Lukaszewicz; Jane Muret; Nadia Smail

SFAR clinical guidelines committee

Alice Blet (President), Marie-Pierre Bonnet (Board representative), Hélène Charbonneau (Secretary), Aurélien Bonnal, Anaïs Caillard, Isabelle Constant (Board representative), Hugues de Courson, Matthieu, Dumont, Denis Frasca, Pierre Huet, El Mahdi Halfiani, Elise Langouet, Axel Maurice-Szamburski, Daphné Michelet, Maxime Nguyen, Stéphanie Ruiz, Michaël Vourc'h.

ANARLF clinical guidelines committee

Thomas Geeraerts (President), Vincent Degos (General secretary), Claire Dahyot-Fizelier (Treasurer), Russel Chabanne, David Couret, Hervé Quintard, Jean-François Payen.

Declaration of competing interest

The authors declare having no conflicts of interest.

Appendices

Glasgow Outcome Scale Extended (GOSE) [15].

GOSE	Poor neurological outcomes
1 Death	
2 Vegetative State	
3 Lower Severe Disability	Totally dependent
4 Upper Severe Disability	Very dependent (but can be left alone several hours)
	Good neurological outcomes
5 Lower Moderate Disability	Partially dependent

(continued on next page)

(continued)

6 Upper Moderate Disability	Partial return to work (or school)
7 Lower Good Recovery	Mild sequelae
8 Upper Good Recovery	Complete recovery

Modified Frailty Index 5 (mFI-5) [29].

Items		Result	Interpretation
Functional dependency	+1	0	Non frail
History of diabetes	+1	1	Pre-frail
Chronic obstructive pulmonary disease	+1	≥2	Frail
Congestive heart failure within 30 days	+1		
Hypertension	+1		

Clinical Frailty Scale (CFS) [31].

Score	Category	Description
1	Very Fit	People who are robust, active, energetic, and motivated. These individuals typically exercise regularly and are among the fittest for their age.
2	Well	People who have no active disease symptoms but are less fit than category 1. They often exercise or are very active occasionally, e.g., seasonally.
3	Managing Well	People whose medical problems are well controlled, but are not regularly active beyond routine walking.
4	Vulnerable	While not dependent on others for daily help, often symptoms limit activities. They often complain of being "slowed up" or have disease symptoms.
5	Mildly Frail	People with limited dependence on others for instrumental activities of daily living (IADLs), such as shopping, transportation, and managing medications.
6	Moderately Frail	People who need help with all outside activities and with keeping house. They need help with both instrumental and non-instrumental activities of daily living (ADLs).
7	Severely Frail	People who are completely dependent on personal care, for most activities of daily living, from physical causes.
8	Very Severely Frail	People who are completely dependent, and can be difficult to engage or frequently seem unaware of their surroundings. They approach the end of life.
9	Terminally Ill	People with a life expectancy of less than six months, who are not otherwise evidently frail.

SPIN score (Penetrating brain injury severity criteria) [226].


Survival score component	Points
motor Glasgow Coma Score	
5-Jan	0
6	9
Pupils	
Non-reactive bilaterally	0
Unequal	4
Globe rupture/non-visual	6
Equal and reactive bilaterally	9
Self-inflicted	
Yes	0
No	4
Transferred	
Yes	0
No	4
Sex	
Male	0
Female	4
Injury Severity Score (ISS)	
≥56	0
41–55	1
25–40	5
≤24	10
INR	
≥2,1	0
1,4–2	6
≤1,3	12

SPIN score ≥ 35 : 98% survival. SPIN score ≤ 20 : 3% survival.

CT scan criteria for poor prognosis in penetrating traumatic brain injury [218,227].



- Multilobar or bihemispheric injury (except bi-frontal trajectory)
- Ventricular injury with hemorrhage
- Diffuse fragmentation
- Missile passing through the geographic center of the brain (i.e., involving the thalamus and basal ganglia)
An area 4 cm above the dorsum sellae was described as the *zona fatalis*
- Trajectory crossing the x, y and z planes
- Midline shift (>10 mm on CT)
- Compressed or obliterated basal cisterns
- Large intracerebral hemorrhage
- Subarachnoid hemorrhage
- Large volume of contused brain
- Posterior fossa wound with brainstem involvement
- 'Tram track sign' hemorrhage on either side of a dark center track in a perforating injury

Figure. Infographic synthesis





NEUROSURGICAL MANAGEMENT OF ADULT AND PEDIATRIC TRAUMATIC BRAIN INJURY DURING THE INITIAL PHASE - Guidelines from the French Society of Neurosurgery 2025




LEVEL OF RECOMMENDATIONS Grade 2 Expert opinion

CTA: Computed tomography angiography; DSA: Digital subtraction angiography; EAM: external auditory meatus; ELD: external lumbar drainage; GCS: Glasgow coma score; ICP: Intracranial pressure; ICHT: intracranial hypertension; MLS: midline shift; pTBI: penetrating traumatic brain injury SDH: subdural hematoma; TBI: traumatic brain injury; TCD: transcranial doppler


Infographics : R. Momet



PRISES EN CHARGE NEUROCHIRURGICALES DES TRAUMATISMES CRANIO-ENCEPHALIQUES A LA PHASE INITIALE RECOMMANDATIONS DE PRATIQUES PROFESSIONNELLES - SFNC 2025



PRISES EN CHARGE NEUROCHIRURGICALES DES TRAUMATISMES CRANIO-ENCEPHALIQUES A LA PHASE INITIALE RECOMMANDATIONS DE PRATIQUES PROFESSIONNELLES - SFNC 2025



Infographie : Z. Monet

References

- Taylor CA, Bell JM, Breiding MJ, Xu L. Traumatic brain injury-related emergency department visits, hospitalizations, and deaths - United States, 2007 and 2013. *MMWR Surveill Summ* 2017;66:1-16. <https://doi.org/10.15585/mmwr.s6609a1>.
- Majdan M, Plancikova D, Brazinova A, Rusnak M, Nieboer D, Feigin V, et al. Epidemiology of traumatic brain injuries in Europe: a cross-sectional analysis. *Lancet Public Health* 2016;1:e76-83. [https://doi.org/10.1016/S2468-2667\(16\)30017-2](https://doi.org/10.1016/S2468-2667(16)30017-2).
- Brazinova A, Rehorkcova V, Taylor MS, Buckova V, Majdan M, Psota M, et al. Epidemiology of traumatic brain injury in Europe: a living systematic review. *J Neurotrauma* 2021;38:1411-40. <https://doi.org/10.1089/neu.2015.4126>.
- Gil-Jardiné C, Payen J-F, Bernard R, Bobbia X, Bouzat P, Catoire P, et al. Management of patients suffering from mild traumatic brain injury 2023. *Anaesth Crit Care Pain Med* 2023;42:101260. <https://doi.org/10.1016/j.accpm.2023.101260>.
- Lumba-Brown A, Yeates KO, Sarmiento K, Breiding MJ, Haegerich TM, Gioia GA, et al. Centers for disease control and prevention guideline on the diagnosis and management of mild traumatic brain injury among children. *JAMA Pediatr* 2018;172:e182853. <https://doi.org/10.1001/jamapediatrics.2018.2853>.
- Carney N, Totten AM, O'Reilly C, Ullman JS, Hawryluk GWJ, Bell MJ, et al. Guidelines for the management of severe traumatic brain injury, fourth edition. *Neurosurgery* 2017;80:6-15. <https://doi.org/10.1227/NEU.0000000000001432>.
- Geeraerts T, Velly L, Abdenour L, Asehnoune K, Audibert G, Bouzat P, et al. Management of severe traumatic brain injury (first 24hours). *Anaesth Crit Care Pain Med* 2018;37:171-86. <https://doi.org/10.1016/j.accpm.2017.12.001>.
- Kochanek PM, Tasker RC, Carney N, Totten AM, Adelson PD, Selden NR, et al. Guidelines for the management of pediatric severe traumatic brain injury, third edition: update of the brain trauma foundation guidelines. *Pediatr Crit Care Med* 2019;20:S1-82. <https://doi.org/10.1097/PCC.0000000000001735>.
- Badjatia N, Carney N, Crocco TJ, Fallat ME, Hennes HMA, Jagoda AS, et al. Guidelines for prehospital management of traumatic brain injury 2nd edition. *Prehosp Emerg Care* 2008;12(Suppl 1):S1-S2. <https://doi.org/10.1080/10903120701732052>.
- Chesnut R, Aguilera S, Buki A, Bulger E, Citerio G, Cooper DJ, et al. A management algorithm for adult patients with both Brain oxygen and intracranial pressure monitoring: the Seattle International Severe Traumatic Brain Injury Consensus Conference (SIBICC). *Intensive Care Med* 2020;46:919-29. <https://doi.org/10.1007/s00134-019-05900-x>.
- Hawryluk GWJ, Aguilera S, Buki A, Bulger E, Citerio G, Cooper DJ, et al. A management algorithm for patients with intracranial pressure monitoring: the Seattle International Severe Traumatic Brain Injury Consensus Conference (SIBICC). *Intensive Care Med* 2019;45:1783-94. <https://doi.org/10.1007/s00134-019-05805-9>.
- Muehlschlegel S, Rajajee V, Wartenberg KE, Alexander SA, Busl KM, Creutzfeldt CJ, et al. Guidelines for neuroprognostication in critically ill adults with moderate-severe traumatic brain injury. *Neurocrit Care* 2024;40:448-76. <https://doi.org/10.1007/s12028-023-01902-2>.
- Bullock MR, Chesnut R, Ghajar J, Gordon D, Hartl R, Newell DW, et al. Surgical management of traumatic parenchymal lesions. *Neurosurgery* 2006;58:S25-46. <https://doi.org/10.1227/01.NEU.0000210365.36914.E3>. discussion S-i-v.
- Hawryluk GWJ, Rubiano AM, Totten AM, O'Reilly C, Ullman JS, Bratton SL, et al. Guidelines for the management of severe traumatic brain injury: 2020 update of the decompressive craniectomy recommendations. *Neurosurgery* 2020;87:427-34. <https://doi.org/10.1093/neuros/nyaa278>.
- https://www.center-tbi.eu/files/approved-translations/French/FRENCH_GOSE.pdf. Accessed July 19th 2024. n.d.
- McCauley SR, Wilde EA, Anderson VA, Bedell G, Beers SR, Campbell TF, et al. Recommendations for the use of common outcome measures in pediatric traumatic brain injury research. *J Neurotrauma* 2012;29:678-705. <https://doi.org/10.1089/neu.2011.1838>.
- Turgeon AF, Lauzier F, Simard J-F, Scales DC, Burns KEA, Moore L, et al. Mortality associated with withdrawal of life-sustaining therapy for patients with severe traumatic brain injury: a Canadian multicenter cohort study. *CMAJ* 2011;183:1581-8. <https://doi.org/10.1503/cmaj.101786>.
- Sahuquillo J, Dennis JA. Decompressive craniectomy for the treatment of high intracranial pressure in closed traumatic brain injury. *Cochrane Database Syst Rev* 2019;12:CD003983. <https://doi.org/10.1002/14651858.CD003983.pub3>.
- Maas AIR, Menon DK, Manley GT, Abrams M, Åkerlund C, Andelic N, et al. Traumatic brain injury: progress and challenges in prevention, clinical care, and

- research. *Lancet Neurol* 2022;21:1004–60. [https://doi.org/10.1016/S1474-4422\(22\)00309-X](https://doi.org/10.1016/S1474-4422(22)00309-X).
- [20] Rohaut B, Calligaris C, Hermann B, Perez P, Faugeras F, Raimondo F, et al. Multimodal assessment improves neuroprognosis performance in clinically unresponsive critical-care patients with brain injury. *Nat Med* 2024. <https://doi.org/10.1038/s41591-024-03019-1>.
 - [21] Retel Helmrich IRA, van Klaveren D, Andelic N, Lingsma H, Maas A, Menon D, et al. Discrepancy between disability and reported well-being after traumatic brain injury. *J Neurol Neurosurg Psychiatry* 2022;93:785–96. <https://doi.org/10.1136/jnnp-2021-326615>.
 - [22] Hirschi R, Rommel C, Letsinger J, Nirula R, Hawryluk GWJ. Brain Trauma Foundation Guideline Compliance: Results of a Multidisciplinary, International Survey. *World Neurosurg* 2018;116:e399–405. <https://doi.org/10.1016/j.wneu.2018.04.215>. Epub 2018 May 9. PMID: 29751187.
 - [23] LOI n° 2016-87 du 2 février 2016 créant de nouveaux droits en faveur des malades et des personnes en fin de vie. <https://www.legifrance.gouv.fr/jorf/id/JORFTEXT000031970253> (accès le 01/07/2024). n.d.
 - [24] Limitations et arrêts des traitements de suppléance vitale chez l'adulte dans le contexte de l'urgence. Recommendations SFMU, SRLF, 2018. https://www.sfmur.org/upload/consensus/RPC_LATenurgencedef_2018.pdf (accès le 01/07/2024). n.d.
 - [25] Souter MJ, Blissitt PA, Blosser S, Bonomo J, Greer D, Jichici D, et al. Recommendations for the critical care management of devastating brain injury: prognostication, psychosocial, and ethical management: a position statement for healthcare professionals from the neurocritical care society. *Neurocrit Care* 2015;23:4–13. <https://doi.org/10.1007/s12028-015-0137-6>.
 - [26] Harvey D, Butler J, Groves J, Manara A, Menon D, Thomas E, et al. Management of perceived devastating brain injury after hospital admission: a consensus statement from stakeholder professional organizations. *Br J Anaesth* 2018;120:138–45. <https://doi.org/10.1016/j.bja.2017.10.002>.
 - [27] Clegg A, Young J, Iliffe S, Rikkert MO, Rockwood K. Frailty in elderly people. *Lancet* 2013;381:752–62. [https://doi.org/10.1016/S0140-6736\(12\)62167-9](https://doi.org/10.1016/S0140-6736(12)62167-9).
 - [28] Velanovich V, Antoine H, Swartz A, Peters D, Rubinfeld I. Accumulating deficits model of frailty and postoperative mortality and morbidity: its application to a national database. *J Surg Res* 2013;183:104–10. <https://doi.org/10.1016/j.jss.2013.01.021>.
 - [29] Subramaniam S, Aalberg JJ, Soriano RP, Divino CM. New 5-factor modified frailty index using American college of surgeons NSQIP data. *J Am Coll Surg* 2018;226:173–181.e8. <https://doi.org/10.1016/j.jamcollsurg.2017.11.005>.
 - [30] Kessler RA, De la Garza Ramos R, Purvis TE, Ahmed AK, Goodwin CR, Sciubba DM, et al. Impact of frailty on complications in patients with thoracic and thoracolumbar spinal fracture. *Clin Neurol Neurosurg* 2018;169:161–5. <https://doi.org/10.1016/j.clineuro.2018.04.014>.
 - [31] Rockwood K, Song X, MacKnight C, Bergman H, Hogan DB, McDowell I, et al. A global clinical measure of fitness and frailty in elderly people. *CMAJ* 2005;173:489–95. <https://doi.org/10.1503/cmaj.050051>.
 - [32] Raïche M, Hébert R, Dubois M-F. PRISMA-7: a case-finding tool to identify older adults with moderate to severe disabilities. *Arch Gerontol Geriatr* 2008;47:9–18. <https://doi.org/10.1016/j.archger.2007.06.004>.
 - [33] Joseph B, Saljuqi AT, Amos JD, Teichman A, Whitmill ML, Anand T, et al. Prospective validation and application of the Trauma-Specific Frailty Index: results of an American Association for the Surgery of Trauma multi-institutional observational trial. *J Trauma Acute Care Surg* 2023;94:36–44. <https://doi.org/10.1097/TA.0000000000003817>.
 - [34] Hoogendijk EO, Afilalo J, Ensrud KE, Kowal P, Onder G, Fried LP. Frailty: implications for clinical practice and public health. *Lancet* 2019;394:1365–75. [https://doi.org/10.1016/S0140-6736\(19\)31786-6](https://doi.org/10.1016/S0140-6736(19)31786-6).
 - [35] Joseph B, Pandit V, Zangbar B, Kulvatunyou N, Hashmi A, Green DJ, et al. Superiority of frailty over age in predicting outcomes among geriatric trauma patients: a prospective analysis. *JAMA Surg* 2014;149:766–72. <https://doi.org/10.1001/jamasurg.2014.296>.
 - [36] Galimberti S, Graziano F, Maas AIR, Isernia G, Lecky F, Jain S, et al. Effect of frailty on 6-month outcome after traumatic brain injury: a multicentre cohort study with external validation. *Lancet Neurol* 2022;21:153–62. [https://doi.org/10.1016/S1474-4422\(21\)00374-4](https://doi.org/10.1016/S1474-4422(21)00374-4).
 - [37] Tang OY, Shao B, Kimata AR, Sastry RA, Wu J, Asaad WF. The impact of frailty on traumatic brain injury outcomes: an analysis of 691 821 nationwide cases. *Neurosurgery* 2022;91:808–20. <https://doi.org/10.1227/neu.0000000000002116>.
 - [38] Abdulle AE, de Koning ME, van der Horn HJ, Scheenen ME, Roks G, Hageman G, et al. Early predictors for long-term functional outcome after mild traumatic brain injury in frail elderly patients. *J Head Trauma Rehabil* 2018;33:E59–67. <https://doi.org/10.1097/HTR.0000000000000368>.
 - [39] Elsamadicy AA, Sandhu MRS, Freedman IG, Koo AB, Reeves BC, Yu J, et al. Impact of frailty on morbidity and mortality in adult patients undergoing surgical evacuation of acute traumatic subdural hematoma. *World Neurosurg* 2022;162:e251–63. <https://doi.org/10.1016/j.wneu.2022.02.122>.
 - [40] Lee H, Tan C, Tran V, Mathew J, Fitzgerald M, Leong R, et al. The utility of the modified frailty index in outcome prediction for elderly patients with acute traumatic subdural hematoma. *J Neurotrauma* 2020;37:2499–506. <https://doi.org/10.1089/neu.2019.6943>.
 - [41] Sastry RA, Feler JR, Shao B, Ali R, McNicoll L, Telfeian AE, et al. Frailty independently predicts unfavorable discharge in non-operative traumatic brain injury: a retrospective single-institution cohort study. *PLoS One* 2022;17:e0275677. <https://doi.org/10.1371/journal.pone.0275677>.
 - [42] Tracy BM, Carlin MN, Tyson JW, Schenker ML, Gelbard RB. The 11-item modified frailty index as a tool to predict unplanned events in traumatic brain injury. *Am Surg* 2020;86:1596–601. <https://doi.org/10.1177/0003134820942196>.
 - [43] Cole KL, Kurudza E, Rahman M, Kazim SF, Schmidt MH, Bowers CA, et al. Use of the 5-factor modified frailty index to predict hospital-acquired infections and length of stay among neurotrauma patients undergoing emergent craniotomy/craniectomy. *World Neurosurg* 2022;164:e1143–52. <https://doi.org/10.1016/j.wneu.2022.05.122>.
 - [44] Herklots MW, Kroon M, Roks G, Oldenbeuving A, Schoonman GG. Poor outcome in frail elderly patient after severe TBI. *Brain Inj* 2022;36:1118–22. <https://doi.org/10.1080/02699052.2022.2109731>.
 - [45] Maragkos GA, Matsoukas S, Cho LD, Legome EL, Wedderburn RV, Margetis K. Comparison of frailty indices and the charlson comorbidity index in traumatic brain injury. *J Head Trauma Rehabil* 2023;38:E177–85. <https://doi.org/10.1097/HTR.0000000000000832>.
 - [46] Rawanduzay C, McIntyre MK, Afridi A, Honig J, Halabi M, Hehir J, et al. The effect of frailty and patient comorbidities on outcomes after acute subdural hemorrhage: a preliminary analysis. *World Neurosurg* 2020;143:e285–93. <https://doi.org/10.1016/j.wneu.2020.07.106>.
 - [47] Yamamoto Y, Hori S, Ushida K, Shirai Y, Shimizu M, Kato Y, et al. Impact of frailty risk on adverse outcomes after traumatic brain injury: a historical cohort study. *J Clin Med* 2022;11:7064. <https://doi.org/10.3390/jcm11237064>.
 - [48] Zacchetti L, Longhi L, Zangari R, Aresi S, Marchesi F, Gritti P, et al. Clinical frailty scale as a predictor of outcome in elderly patients affected by moderate or severe traumatic brain injury. *Front Neurol* 2023;14:1021020. <https://doi.org/10.3389/fneur.2023.1021020>.
 - [49] Roohollahi F, Molavi S, Mohammadi M, Mohamadi M, Mohammadi A, Kankam SB, et al. Prognostic value of frailty for outcome following traumatic brain injury: a systematic review and meta-analysis. *J Neurotrauma* 2024;41:331–48. <https://doi.org/10.1089/neu.2023.0176>.
 - [50] MRC CRASH Trial Collaborators, Perel P, Arango M, Clayton T, Edwards P, Komolafe E, et al. Predicting outcome after traumatic brain injury: practical prognostic models based on large cohort of international patients. *BMJ* 2008;336:425–9. <https://doi.org/10.1136/bmj.39461.643438.25>.
 - [51] Murray GD, Butcher I, McHugh GS, Lu J, Mushkudiani NA, Maas AIR, et al. Multivariable prognostic analysis in traumatic brain injury: results from the IMPACT study. *J Neurotrauma* 2007;24:329–37. <https://doi.org/10.1089/neu.2006.0035>.
 - [52] Pancykowski DM, Puccio AM, Scruggs BJ, Bauer JS, Hricik AJ, Beers SR, et al. Prospective independent validation of IMPACT modeling as a prognostic tool in severe traumatic brain injury. *J Neurotrauma* 2012;29:47–52. <https://doi.org/10.1089/neu.2010.1482>.
 - [53] Timmons SD, Bee T, Webb S, Diaz-Arrastia RR, Hesdorffer D. Using the abbreviated injury severity and Glasgow Coma Scale scores to predict 2-week mortality after traumatic brain injury. *J Trauma* 2011;71:1172–8. <https://doi.org/10.1097/TA.0b013e31822b0f4b>.
 - [54] Steyerberg EW, Mushkudiani N, Perel P, Butcher I, Lu J, McHugh GS, et al. Predicting outcome after traumatic brain injury: development and international validation of prognostic scores based on admission characteristics. *PLoS Med* 2008;5:e165. <https://doi.org/10.1371/journal.pmed.0050165>. discussion e165.
 - [55] Brennan PM, Murray GD, Teasdale GM. Simplifying the use of prognostic information in traumatic brain injury. Part 1: the GCS-Pupils score: an extended index of clinical severity. *J Neurosurg* 2018;128:1612–20. <https://doi.org/10.3171/2017.12.JNS172780>.
 - [56] Gao G, Wu X, Feng J, Hui J, Mao Q, Lecky F, et al. Clinical characteristics and outcomes in patients with traumatic brain injury in China: a prospective, multicentre, longitudinal, observational study. *Lancet Neurol* 2020;19:670–7. [https://doi.org/10.1016/S1474-4422\(20\)30182-4](https://doi.org/10.1016/S1474-4422(20)30182-4).
 - [57] Ziaierad M, Alimohammadi N, Trajipour A, Aminmansour B. Association between outcome of severe traumatic brain injury and demographic, clinical, injury-related variables of patients. *Iran J Nurs Midwifery Res* 2018;23:211–6. <https://doi.org/10.4103/ijnmr.IJNMR.65.17>.
 - [58] Chiericato A, Venditto A, Russo E, Martino C, Bini G. Aggressive medical management of acute traumatic subdural hematomas before emergency craniotomy in patients presenting with bilateral unreactive pupils. A cohort study. *Acta Neurochir (Wien)* 2017;159:1553–9. <https://doi.org/10.1007/s00701-017-3190-4>.
 - [59] Lan Z, Richard SA, Li Q, Wu C, Zhang Q, Chen R, et al. Outcomes of patients undergoing craniotomy and decompressive craniectomy for severe traumatic brain injury with brain herniation: a retrospective study. *Medicine (Baltimore)* 2020;99:e22742. <https://doi.org/10.1097/MD.00000000000022742>.
 - [60] Yue JK, Lee YM, Sun X, van Essen TA, Elguindy MM, Belton PJ, et al. Performance of the IMPACT and CRASH prognostic models for traumatic brain injury in a contemporary multicenter cohort: a TRACK-TBI study. *J Neurosurg* 2024;141:417–29. <https://doi.org/10.3171/2023.11.JNS231425>.
 - [61] Scotter J, Hendrickson S, Marcus HJ, Wilson MH. Prognosis of patients with bilateral fixed dilated pupils secondary to traumatic extradural or subdural haematoma who undergo surgery: a systematic review and meta-analysis. *Emerg Med J* 2015;32:654–9. <https://doi.org/10.1136/emered-2014-204260>.
 - [62] Griep DW, Miller A, Sorek S, Rahme R. Are bilaterally fixed and dilated pupils the kiss of death in patients with transtentorial herniation? Systematic review and pooled analysis. *World Neurosurg* 2022;164:e427–35. <https://doi.org/10.1016/j.wneu.2022.04.118>.
 - [63] Griep DW, Miller A, Sorek S, Moawad S, Rahme R. Bilaterally fixed and dilated pupils are not the kiss of death in patients with transtentorial herniation: a single surgeon's experience. *World Neurosurg* 2022;167:e444–50. <https://doi.org/10.1016/j.wneu.2022.08.036>.

- [64] Griep DW, Miller A, Sorek S, Naeem K, Moawad S, Klein D, et al. Emergency decompressive surgery in patients with transtentorial brain herniation and pupillary abnormalities: the importance of improved pupillary response after osmotherapy and surgery. *J Neurosurg* 2024;140:544–51. <https://doi.org/10.3171/2023.5.JNS23163>.
- [65] Cohen JE, Montero A, Israel ZH. Prognosis and clinical relevance of anisocoria-craniotomy latency for epidural hematoma in comatose patients. *J Trauma* 1996;41:120–2. <https://doi.org/10.1097/00005373-199607000-00019>.
- [66] Dent DL, Croce MA, Menke PG, Young BH, Hinson MS, Kudsk KA, et al. Prognostic factors after acute subdural hematoma. *J Trauma* 1995;39:36–42. <https://doi.org/10.1097/00005373-199507000-00005>. discussion 42–43.
- [67] Haselsberger K, Pucher R, Auer LM. Prognosis after acute subdural or epidural haemorrhage. *Acta Neurochir (Wien)* 1988;90:111–6. <https://doi.org/10.1007/BF01560563>.
- [68] Joosse P, Saltzherr T-P, van Lieshout WAM, van Exter P, Ponsen K-J, Vandertop WP, et al. Impact of secondary transfer on patients with severe traumatic brain injury. *J Trauma Acute Care Surg* 2012;72:487–90. <https://doi.org/10.1097/TA.0b013e318226ed59>.
- [69] Matsushima K, Inaba K, Siboni S, Skiada D, Strumwasser AM, Magee GA, et al. Emergent operation for isolated severe traumatic brain injury: does time matter? *J Trauma Acute Care Surg* 2015;79:838–42. <https://doi.org/10.1097/TA.0000000000000719>.
- [70] Mendelow AD, Karmi MZ, Paul KS, Fuller GA, Gillingham FJ. Extradural haematoma: effect of delayed treatment. *Br Med J* 1979;1:1240–2. <https://doi.org/10.1136/bmj.1.6173.1240>.
- [71] Seelig JM, Becker DP, Miller JD, Greenberg RP, Ward JD, Choi SC. Traumatic acute subdural hematoma: major mortality reduction in comatose patients treated within four hours. *N Engl J Med* 1981;304:1511–8. <https://doi.org/10.1056/NEJM198106183042503>.
- [72] Tien HCN, Jung V, Pinto R, Mainprize T, Scales DC, Rizoli SB. Reducing time-to-treatment decreases mortality of trauma patients with acute subdural hematoma. *Ann Surg* 2011;253:1178–83. <https://doi.org/10.1097/SLA.0b013e318217e339>.
- [73] Zafrullah Arifin M, Gunawan W. Analysis of presurgery time as a prognostic factor in traumatic acute subdural hematoma. *J Neurosurg Sci* 2013;57:277–80.
- [74] Gupta S, Khajanchi M, Kumar V, Raykar NP, Alkire BC, Roy N, et al. Third delay in traumatic brain injury: time to management as a predictor of mortality. *J Neurosurg* 2020;132:289–95. <https://doi.org/10.3171/2018.8.JNS182182>.
- [75] Harmsen AMK, Giannakopoulos GF, Moerbeek PR, Jansma EP, Bonjer HJ, Bloemers FW. The influence of prehospital time on trauma patients outcome: a systematic review. *Injury* 2015;46:602–9. <https://doi.org/10.1016/j.injury.2015.01.008>.
- [76] Kim YJ. The impact of time from ED arrival to surgery on mortality and hospital length of stay in patients with traumatic brain injury. *J Emerg Nurs* 2011;37:328–33. <https://doi.org/10.1016/j.jen.2010.04.017>.
- [77] Shackelford SA, Del Junco DJ, Reade MC, Bell R, Becker T, Gurney J, et al. Association of time to craniotomy with survival in patients with severe combat-related brain injury. *Neurosurg Focus* 2018;45:E2. <https://doi.org/10.3171/2018.9.FOCUS18404>.
- [78] Vaca SD, Kuo BJ, Nickenig Vissoci JR, Staton CA, Xu LW, Muhumuza M, et al. Temporal delays along the neurosurgical care continuum for traumatic brain injury patients at a tertiary care hospital in Kampala, Uganda. *Neurosurgery* 2019;84:95–103. <https://doi.org/10.1093/neuros/nyy004>.
- [79] De Vloot P, Nijs S, Verelst S, van Loon J, Depreitere B. Prehospital and intrahospital temporal intervals in patients requiring emergent trauma craniotomy. A 6-year observational study in a level 1 trauma center. *World Neurosurg* 2018;114:e546–58. <https://doi.org/10.1016/j.wneu.2018.03.032>.
- [80] Bartlett JR, Neil-Dwyer G. Extradural haematoma: effect of delayed treatment. *Br Med J* 1979;2:440–1. <https://doi.org/10.1136/bmj.2.6187.440-b>.
- [81] Bonow RH, Barber J, Temkin NR, Videtta W, Rondina C, Petroni G, et al. The Outcome of severe traumatic brain injury in Latin America. *World Neurosurg* 2018;111:e82–90. <https://doi.org/10.1016/j.wneu.2017.11.171>.
- [82] Evenden J, Harris D, Wells AJ, Toson B, Ellis DY, Lambert PF. Increased distance or time from a major trauma centre in South Australia is not associated with worse outcomes after moderate to severe traumatic brain injury. *Emerg Med Australas* 2023;35:998–1004. <https://doi.org/10.1111/1742-6723.14281>.
- [83] Hu J, Sokh V, Nguon S, Heng YV, Husum H, Kloster R, et al. Emergency craniotomy and burr-hole trephination in a low-resource setting: capacity building at a regional hospital in Cambodia. *Int J Environ Res Public Health* 2022;19:6471. <https://doi.org/10.3390/ijerph19116471>.
- [84] Mejjaddam AY, Elmer J, Sideris AC, Chang Y, Petrovick L, Alam HB, et al. Prolonged emergency department length of stay is not associated with worse outcomes in traumatic brain injury. *J Emerg Med* 2013;45:384–91. <https://doi.org/10.1016/j.jemermed.2013.04.015>.
- [85] Walcott BP, Khanna A, Kwon C-S, Phillips HW, Nahed BV, Coumans J-V. Time interval to surgery and outcomes following the surgical treatment of acute traumatic subdural hematoma. *J Clin Neurosci* 2014;21:2107–11. <https://doi.org/10.1016/j.jocn.2014.05.016>.
- [86] Tonui PM, Spilman SK, Pelaez CA, Mankins MR, Sidwell RA. Head CT before transfer does not decrease time to craniotomy for TBI patients. *Am Surg* 2018;84:201–7.
- [87] Vats A, Roy D, Prasad MK. Direct versus indirect transfer for traumatic brain injury to James Cook University Hospital: a retrospective study. *Ann R Coll Surg Engl* 2021;103:23–8. <https://doi.org/10.1308/rcsann.2020.0180>.
- [88] Sergides IG, Whiting G, Howarth S, Hutchinson PJ. Is the recommended target of 4 hours from head injury to emergency craniotomy achievable? *Br J Neurosurg* 2006;20:301–5. <https://doi.org/10.1080/02688690600999976>.
- [89] Wilberger JE, Harris M, Diamond DL. Acute subdural hematoma: morbidity, mortality, and operative timing. *J Neurosurg* 1991;74. <https://doi.org/10.3171/jns.1991.74.2.0212>.
- [90] Kotwicz Z, Brzeziński J. Acute subdural haematoma in adults: an analysis of outcome in comatose patients. *Acta Neurochir (Wien)* 1993;121:95–9. <https://doi.org/10.1007/BF01809257>.
- [91] Fountain DM, Kolias AG, Lecky FE, Bouamra O, Lawrence T, Adams H, et al. Survival trends after surgery for acute subdural hematoma in adults over a 20-year period. *Ann Surg* 2017;265:590–6. <https://doi.org/10.1097/SLA.0000000000001682>.
- [92] Gale SC, Peters J, Hansen A, Dombrovskiy VY, Detwiler PW. Impact of transfer distance and time on rural brain injury outcomes. *Brain Inj* 2016;30:437–40. <https://doi.org/10.3109/02699052.2016.1140808>.
- [93] Schellenberg M, Benjamin E, Owattanapanich N, Inaba K, Demetriades D. The impact of delayed time to first CT head in traumatic brain injury. *Eur J Trauma Emerg Surg* 2021;47:1511–6. <https://doi.org/10.1007/s00068-020-01421-1>.
- [94] Tohme S, Delhumeau C, Zuercher M, Haller G, Walder B. Prehospital risk factors of mortality and impaired consciousness after severe traumatic brain injury: an epidemiological study. *Scand J Trauma Resusc Emerg Med* 2014;22:1. <https://doi.org/10.1186/1757-7241-22-1>.
- [95] Trivedi DJ, Bass GA, Forssten MP, Scheufler K-M, Olivecrona M, Cao Y, et al. The significance of direct transportation to a trauma center on survival for severe traumatic brain injury. *Eur J Trauma Emerg Surg* 2022;48:2803–11. <https://doi.org/10.1007/s00068-022-01885-3>.
- [96] Zimmerman A, Fox S, Griffin R, Nelp T, Thomaz EBAF, Mvungi M, et al. An analysis of emergency care delays experienced by traumatic brain injury patients presenting to a regional referral hospital in a low-income country. *PLoS One* 2020;15:e0240528. <https://doi.org/10.1371/journal.pone.0240528>.
- [97] Newgard CD, Meier EN, Bulger EM, Buick J, Sheehan K, Lin S, et al. Revisiting the “Golden Hour”: an evaluation of out-of-hospital time in shock and traumatic brain injury. *Ann Emerg Med* 2015;66:30–41.e3. <https://doi.org/10.1016/j.annemergmed.2014.12.004>.
- [98] Sashida R, Suehiro E, Tanaka T, Shiomi N, Yatsushige H, Hirota S, et al. Effects of dual antiplatelet therapy (DAPT) compared to single antiplatelet therapy (SAPT) in patients with traumatic brain injury. *Clin Neurol Neurosurg* 2024;244:108418. <https://doi.org/10.1016/j.clineuro.2024.108418>.
- [99] Yorkgitis BK, Tatum DM, Taghavi S, Schroepel TJ, Noorbakhsh MR, Philips FH, et al. Eastern Association for the Surgery of Trauma Multicenter Trial: comparison of pre-injury antithrombotic use and reversal strategies among severe Traumatic brain injury patients. *J Trauma Acute Care Surg* 2022;92:88–92. <https://doi.org/10.1097/TA.00000000000003421>.
- [100] Cheng L, Cui G, Yang R. The impact of preinjury use of antiplatelet drugs on outcomes of traumatic brain injury: a systematic review and meta-analysis. *Front Neurol* 2022;13. <https://doi.org/10.3389/fneur.2022.724641>.
- [101] Greuter L, Ullmann M, Mariani L, Guzman R, Soleman J. Effect of preoperative antiplatelet or anticoagulation therapy on hemorrhagic complications in patients with traumatic brain injury undergoing craniotomy or craniectomy. *Neurosurg Focus* 2019;47:E3. <https://doi.org/10.3171/2019.8.FOCUS19546>.
- [102] Han H, Koh EJ, Choi H, Kim B-C, Yang SY, Cho K-T. The Effect of preoperative antiplatelet therapy on hemorrhagic complications after decompressive craniectomy in patients with traumatic brain injury. *Korean J Neurotrauma* 2016;12:61. <https://doi.org/10.13004/kjnt.2016.12.2.61>.
- [103] Senft C, Schuster T, Forster M-T, Seifert V, Gerlach R. Management and outcome of patients with acute traumatic subdural hematomas and pre-injury oral anticoagulation therapy. *Neurol Res* 2009;31:1012–8. <https://doi.org/10.1179/174313209X409034>.
- [104] Suehiro E, Fujiyama Y, Kiyohira M, Haji K, Ishihara H, Nomura S, et al. Risk of deterioration of geriatric traumatic brain injury in patients treated with antithrombotic drugs. *World Neurosurg* 2019;127:e1221–7. <https://doi.org/10.1016/j.wneu.2019.04.108>.
- [105] Scotti P, Séguin C, Lo BWY, Guise E de, Troquet J-M, Marcoux J. Antithrombotic agents and traumatic brain injury in the elderly population: hemorrhage patterns and outcomes. *J Neurosurg* 2019;133:486–95. <https://doi.org/10.3171/2019.4.JNS19252>.
- [106] Vehviläinen J, Virta JJ, Skrifvars MB, Reinikainen M, Bendel S, Ala-Kokko T, et al. Effect of antiplatelet and anticoagulant medication use on injury severity and mortality in patients with traumatic brain injury treated in the intensive care unit. *Acta Neurochir* 2023;165:4003–12. <https://doi.org/10.1007/s00701-023-05850-w>.
- [107] Rafieezadeh A, Zangbar B, Zeeshan M, Gandhi C, Al-Mufti F, Jehan F, et al. Predictors of mortality after craniotomy for geriatric traumatic brain injury. *Injury* 2024;55:111585. <https://doi.org/10.1016/j.injury.2024.111585>.
- [108] Hecht JP, LaDuke ZJ, Cain-Nielsen AH, Hemmilla MR, Wahl WL. Effect of pre-injury oral anticoagulants on outcomes following traumatic brain injury from falls in older adults. *Pharmacotherapy* 2020;40:604–13. <https://doi.org/10.1002/phar.2435>.
- [109] Lim XT, Ang E, Lee ZX, Hajibandeh S, Hajibandeh S. Prognostic significance of preinjury anticoagulation in patients with traumatic brain injury: a systematic review and meta-analysis. *J Trauma Acute Care Surg* 2021;90:191. <https://doi.org/10.1097/TA.00000000000002976>.
- [110] Tollefsen MH, Vik A, Skandsen T, Sandrød O, Deane SF, Rao V, et al. Patients with moderate and severe traumatic brain injury: impact of preinjury platelet inhibitor or warfarin treatment. *World Neurosurg* 2018;114:e209–17. <https://doi.org/10.1016/j.wneu.2018.02.167>.
- [111] Dang DD, Mugge LA, Ramanathan P, Dang JV, Awan OK, Diekemper N, et al. The implications of antithrombotic agents on subdural hematoma evacuation: what

- does “reversal” truly entail? *Neurosurg Focus* 2023;55:E5. <https://doi.org/10.3171/2023.7.FOCUS23354>.
- [112] Julien J, Alsieiri G, Marcoux J, Hasen M, Correa JA, Feyz M, et al. Antithrombotic agents intake prior to injury does not affect outcome after a traumatic brain injury in hospitalized elderly patients. *J Clin Neurosci* 2017;38:122–5. <https://doi.org/10.1016/j.jocn.2016.12.032>.
- [113] Prexl O, Bruckbauer M, Voelckel W, Grottko O, Ponschab M, Maegele M, et al. The impact of direct oral anticoagulants in traumatic brain injury patients greater than 60-years-old. *Scand J Trauma Resusc Emerg Med* 2018;26:20. <https://doi.org/10.1186/s13049-018-0487-0>.
- [114] Pang CH, Lee SE, Yoo H. Clinical factors and perioperative strategies associated with outcome in preinjury antiplatelet and anticoagulation therapy for patients with traumatic brain injuries. *J Korean Neurosurg Soc* 2015;58:262–70. <https://doi.org/10.3340/jkns.2015.58.3.262>.
- [115] Bhau KS, Bhau SS, Dhar S, Kachroo SL, Babu ML, Chungoo RK. Traumatic extradural hematoma - role of non-surgical management and reasons for conversion. *Indian J Surg* 2010;72:124–9. <https://doi.org/10.1007/s12262-010-0036-1>.
- [116] Bullock MR, Chesnut R, Ghajar J, Gordon D, Hartl R, Newell DW, et al. Surgical management of acute epidural hematomas. *Neurosurgery* 2006;58:S7–15. discussion Si-iv.
- [117] Gutowski P, Meier U, Rohde V, Lemcke J, von der Brelie C. Clinical outcome of epidural hematoma treated surgically in the era of modern resuscitation and trauma care. *World Neurosurg* 2018;118:e166–74. <https://doi.org/10.1016/j.wneu.2018.06.147>.
- [118] Korean Trauma Data Bank System Committee, Jeong YH, Oh JW, Cho S. Clinical outcome of acute epidural hematoma in Korea: preliminary report of 285 cases registered in the Korean trauma data bank system. *Korean J Neurotrauma* 2016;12:47–54. <https://doi.org/10.13004/kjnt.2016.12.2.47>.
- [119] Kang J, Hong S, Hu C, Pyen J, Whang K, Cho S, et al. Clinical analysis of delayed surgical epidural hematoma. *Korean J Neurotrauma* 2015;11:112–7. <https://doi.org/10.13004/kjnt.2015.11.2.112>.
- [120] Basamh M, Robert A, Lamoureux J, Saluja RS, Marcoux J. Epidural hematoma treated conservatively: when to expect the worst. *Can J Neurol Sci* 2016;43:74–81. <https://doi.org/10.1017/cjn.2015.232>.
- [121] Chen TY, Wong CW, Chang CN, Lui TN, Cheng WC, Tsai MD, et al. The expectant treatment of “asymptomatic” supratentorial epidural hematomas. *Neurosurgery* 1993;32:176–9. <https://doi.org/10.1227/00006123-199302000-00004>. discussion 179.
- [122] Cheung PSY, Lam JMY, Yeung JHH, Graham CA, Rainer TH. Outcome of traumatic extradural haematoma in Hong Kong. *Injury* 2007;38:76–80. <https://doi.org/10.1016/j.injury.2006.08.059>.
- [123] Offner PJ, Pham B, Hawkes A. Nonoperative management of acute epidural hematomas: a “no-brainer.”. *Am J Surg* 2006;192:801–5. <https://doi.org/10.1016/j.amjsurg.2006.08.047>.
- [124] Guo C, Liu L, Wang B, Wang Z. Swirl sign in traumatic acute epidural hematoma: prognostic value and surgical management. *Neurol Sci* 2017;38:2111–6. <https://doi.org/10.1007/s10072-017-3121-4>.
- [125] Beniwal HK, Reddy MS, Rao GP, Srikrishnaditya M, Beniwal A. Prognostic value of swirl sign in acute epidural hemorrhage. *Asian J Neurosurg* 2022;17:584–7. https://doi.org/10.4103/ajns.AJNS_68_21.
- [126] Kvint S, Gutierrez A, Blue R, Petrov D. Surgical management of trauma-related intracranial hemorrhage-A review. *Curr Neurol Neurosci Rep* 2020;20:63. <https://doi.org/10.1007/s11910-020-01080-0>.
- [127] Bullock MR, Chesnut R, Ghajar J, Gordon D, Hartl R, Newell DW, et al. Surgical management of posterior fossa mass lesions. *Neurosurgery* 2006;58:S47–55. <https://doi.org/10.1227/01.NEU.0000210366.36914.38>. discussion Si-iv.
- [128] Sharafat S, Azam F, Khan Z. Posterior fossa epidural hematoma: a 6-year management experience. *Pak J Med Sci* 2023;39:253–6. <https://doi.org/10.12669/pjms.39.1.6408>.
- [129] Daoud SS, Jamous MA, Al Barbarawi MM, Jarrar S, Jaradat A, Aljabali AS, et al. Operative versus non-operative management of posterior fossa epidural hematoma: a systematic review and meta-analysis. *Neurochirurgie* 2024;70:101578. <https://doi.org/10.1016/j.neuchi.2024.101578>.
- [130] Fernandes-Cabral DT, Kooshkabad A, Panesar SS, Celticki E, Borghie-Razavi H, Celticki P, et al. Surgical management of vertex epidural hematoma: technical case report and literature review. *World Neurosurg* 2017;103:475–83. <https://doi.org/10.1016/j.wneu.2017.04.040>.
- [131] Parker SL, Kabani AA, Conner CR, Choi PA, Withrow JS, Cai C, et al. Management of venous sinus-related epidural hematomas. *World Neurosurg* 2020;138:e241–50. <https://doi.org/10.1016/j.wneu.2020.02.089>.
- [132] Oertel M, Kelly DF, McArthur D, Boscardin WJ, Glenn TC, Lee JH, et al. Progressive hemorrhage after head trauma: predictors and consequences of the evolving injury. *J Neurosurg* 2002;96:109–16. <https://doi.org/10.3171/jns.2002.96.1.0109>.
- [133] Bezircioğlu H, Erşahin Y, Demirci F, Yurt I, Dönertaş K, Tektaş S. Nonoperative treatment of acute extradural hematomas: analysis of 80 cases. *J Trauma* 1996;41:696–8. <https://doi.org/10.1097/00005373-199610000-00016>.
- [134] Sullivan TP, Jarvik JG, Cohen WA. Follow-up of conservatively managed epidural hematomas: implications for timing of repeat CT. *AJNR Am J Neuroradiol* 1999;20:107–13.
- [135] Gregori F, Santoro G, Mancarella C, Piccirilli M, Domenicucci M. Development of a delayed acute epidural hematoma following contralateral epidural hematoma evacuation: case report and review of literature. *Acta Neurol Belg* 2019;119:15–20. <https://doi.org/10.1007/s13760-018-1049-y>.
- [136] Bullock MR, Chesnut R, Ghajar J, Gordon D, Hartl R, Newell DW, et al. Surgical management of acute subdural hematomas. *Neurosurgery* 2006;58:S16–24. discussion Si-iv.
- [137] Mathew P, Oluoch-Olunya DL, Condon BR, Bullock R. Acute subdural haematoma in the conscious patient: outcome with initial non-operative management. *Acta Neurochir (Wien)* 1993;121:100–8. <https://doi.org/10.1007/BF01809258>.
- [138] Servadei F, Nasi MT, Cremonini AM, Giuliani G, Cenni P, Nanni A. Importance of a reliable admission Glasgow Coma Scale score for determining the need for evacuation of posttraumatic subdural hematomas: a prospective study of 65 patients. *J Trauma* 1998;44:868–73. <https://doi.org/10.1097/00005373-199805000-00021>.
- [139] Wong CW. Criteria for conservative treatment of supratentorial acute subdural haematomas. *Acta Neurochir (Wien)* 1995;135:38–43. <https://doi.org/10.1007/BF02307412>.
- [140] van Essen TA, Lingsma HF, Piscià D, Singh RD, Volovici V, den Boogert HF, et al. Surgery versus conservative treatment for traumatic acute subdural haematoma: a prospective, multicentre, observational, comparative effectiveness study. *Lancet Neurol* 2022;21:620–31. [https://doi.org/10.1016/S1474-4422\(22\)00166-1](https://doi.org/10.1016/S1474-4422(22)00166-1).
- [141] Karibe H, Hayashi T, Hirano T, Kameyama M, Nakagawa A, Tominaga T. Surgical management of traumatic acute subdural hematoma in adults: a review. *Neurol Med Chir (Tokyo)* 2014;54:887–94. <https://doi.org/10.2176/nmc.cr.2014-0204>.
- [142] Raj R, Mikkonen ED, Kivisaari R, Skrifvars MB, Korja M, Siironen J. Mortality in elderly patients operated for an acute subdural hematoma: a surgical case series. *World Neurosurg* 2016;88:592–7. <https://doi.org/10.1016/j.wneu.2015.10.095>.
- [143] Mak CHK, Wong SKH, Wong GK, Ng S, Wang KKW, Lam PK, et al. Traumatic brain injury in the elderly: is it as bad as we think? *Curr Tran Geriatr Gerontol Rep* 2012;1:171–8. <https://doi.org/10.1007/s13670-012-0017-2>.
- [144] Essen TA van, Erp IAM van, Lingsma HF, Piscià D, Yue JK, Singh RD, et al. Comparative effectiveness of decompressive craniectomy versus craniotomy for traumatic acute subdural hematoma (CENTER-TBI): an observational cohort study. *eClinicalMedicine* 2023;63. <https://doi.org/10.1016/j.eclinm.2023.102161>.
- [145] Gavrilu Laic RA, Sloten JV, Depreitere B. Neurosurgical treatment in elderly patients with Traumatic brain injury: a 20-year follow-up study. *Brain and Spine* 2023;3:101723. <https://doi.org/10.1016/j.bas.2023.101723>.
- [146] Evans LR, Jones J, Lee HQ, Gantner D, Jaison A, Matthew J, et al. Prognosis of acute subdural hematoma in the elderly: a systematic review. *J Neurotrauma* 2019;36:517–22. <https://doi.org/10.1089/neu.2018.5829>.
- [147] Borkar SA, Sinha S, Agrawal D, Satyarthee GD, Gupta D, Mahapatra AK. Severe head injury in the elderly: risk factor assessment and outcome analysis in a series of 100 consecutive patients at a Level 1 trauma centre. *Indian J Neurotrauma* 2011;08:77–82. [https://doi.org/10.1016/S0973-0508\(11\)80004-3](https://doi.org/10.1016/S0973-0508(11)80004-3).
- [148] Cnossen MC, Scholten AC, Lingsma HF, Synnot A, Tavender E, Gantner D, et al. Adherence to guidelines in adult patients with traumatic brain injury: a living systematic review. *J Neurotrauma* 2021;38:1072–85. <https://doi.org/10.1089/neu.2015.4121>.
- [149] Singh RD, van Dijk JTJM, van Essen TA, Lingsma HF, Polinder SS, Kompanje EJO, et al. Randomized Evaluation of Surgery in Elderly with Traumatic Acute SubDural Hematoma (RESET-ASDH trial): study protocol for a pragmatic randomized controlled trial with multicenter parallel group design. *Trials* 2022;23:242. <https://doi.org/10.1186/s13063-022-06184-1>.
- [150] Singh RD, van Dijk JTJM, van Essen TA, Nix HP, Vreeburg RJG, den Boogert HF, et al. The death of a neurotrauma trial lessons learned from the prematurely halted randomized evaluation of surgery in elderly with traumatic acute subdural hematoma (RESET-ASDH) trial. *Brain Spine* 2024;4:102903. <https://doi.org/10.1016/j.bas.2024.102903>.
- [151] Phuenpathom N, Choomuang M, Ratanalert S. Outcome and outcome prediction in acute subdural hematoma. *Surg Neurol* 1993;40:22–5. [https://doi.org/10.1016/0090-3019\(93\)90164-v](https://doi.org/10.1016/0090-3019(93)90164-v).
- [152] Fearnside MR, Cook RJ, McDougall P, McNeil RJ. The Westmead Head Injury Project outcome in severe Head Injury. A comparative analysis of pre-hospital, clinical and CT variables. *Br J Neurosurg* 1993;7:267–79. <https://doi.org/10.3109/02688699309023809>.
- [153] Leitgeb J, Mauritz W, Brazinova A, Janciak I, Majdan M, Wilbacher I, et al. Outcome after severe brain trauma due to acute subdural hematoma. *J Neurosurg* 2012;117:324–33. <https://doi.org/10.3171/2012.4.JNS111448>.
- [154] van Essen TA, Res L, Schoones J, de Ruiter G, Dekkers O, Maas A, et al. Mortality reduction of acute surgery in traumatic acute subdural hematoma since the 19th century: systematic review and meta-analysis with dramatic effect: is surgery the obvious parachute? *J Neurotrauma* 2023;40:22–32. <https://doi.org/10.1089/neu.2022.0137>.
- [155] van Essen TA, Dijkman MD, Cnossen MC, Moudrout W, Ardon H, Schoonman GG, et al. Comparative effectiveness of surgery for traumatic acute subdural hematoma in an aging population. *J Neurotrauma* 2019;36:1184–91. <https://doi.org/10.1089/neu.2018.5869>.
- [156] van Essen TA, de Ruiter GCW, Kho KH, Peul WC. Neurosurgical treatment variation of traumatic brain injury: evaluation of acute subdural hematoma management in Belgium and The Netherlands. *J Neurotrauma* 2017;34:881–9. <https://doi.org/10.1089/neu.2016.4495>.
- [157] Bajsarowicz P, Prakash I, Lamoureux J, Saluja RS, Feyz M, Maleki M, et al. Nonsurgical acute traumatic subdural hematoma: what is the risk? *J Neurosurg* 2015;123:1176–83. <https://doi.org/10.3171/2014.10.JNS141728>.
- [158] Kayahara T, Kikkawa Y, Komine H, Kamide T, Suzuki K, Shibata A, et al. Predictors of subacute hematoma expansion requiring surgical evacuation after initial conservative treatment in patients with acute subdural hematoma. *Acta*

- Neurochir (Wien) 2020;162:357–63. <https://doi.org/10.1007/s00701-019-04187-7>.
- [159] Kashkoush AI, Whiting BB, Desai A, Pettit JC, El-Abtah ME, Mcmillan A, et al. Clinical outcomes after nonoperative management of large acute traumatic subdural hematomas in older patients: a propensity-scored retrospective analysis. *Neurosurgery* 2023;92:293–9. <https://doi.org/10.1227/NEU.0000000000002192>.
- [160] Aziz MM, Molla S, Abdelrahman HA, Dawood OM. Depressed skull fractures overlying dural venous sinuses: management modalities and review of literature. *Turk Neurosurg* 2019;29:856–63. <https://doi.org/10.5137/1019-5149.JTN.25572-18.2>.
- [161] Hussein AA, Laeke T, Yohannes D. Clinical outcomes and prognostic determinants of surgically treated depressed skull fracture in Addis Ababa university neurosurgical teaching hospitals: a prospective multicenter observational study. *World Neurosurg* 2022;158:e55–63. <https://doi.org/10.1016/j.wneu.2021.10.089>.
- [162] Kumar S, Nahar S, Sahana D, Jain AK, Rathore L, Chandrakar N, et al. A prospective comparative evaluation of surgery and conservative treatment for compound depressed skull fractures. *World Neurosurg* 2022;164:e1281–9. <https://doi.org/10.1016/j.wneu.2022.06.019>.
- [163] Satardey RS, Balasubramaniam S, Pandya JS, Mahey RC. Analysis of factors influencing outcome of depressed fracture of skull. *Asian J Neurosurg* 2018;13:341–7. https://doi.org/10.4103/ajns.AJNS_117_16.
- [164] Zapata L, Wright EJ, Nakaji P. Clinical and neurosurgical management of cranial machete injuries: the experience of a tertiary referral center in Nicaragua. *World Neurosurg* 2018;116:140–3. <https://doi.org/10.1016/j.wneu.2018.05.020>.
- [165] Salia SM, Mersha HB, Aklilu AT, Baleh AS, Lund-Johansen M. Predicting dural tear in compound depressed skull fractures: a prospective multicenter correlational study. *World Neurosurg* 2018;114:e833–9. <https://doi.org/10.1016/j.wneu.2018.03.095>.
- [166] Bullock MR, Chesnut R, Ghajar J, Gordon D, Hartl R, Newell DW, et al. Surgical management of depressed cranial fractures. *Neurosurgery* 2006;58:S56–60. <https://doi.org/10.1227/01.NEU.0000210367.14043.0E>. discussion Si-iv.
- [167] Mukherjee KK, Dhandapani S, Sarda AC, Tripathi M, Salunke P, Srinivasan A, et al. Prospective comparison of simple suturing and elevation debridement in compound depressed fractures with no significant mass effect. *Acta Neurochir (Wien)* 2015;157:305–9. <https://doi.org/10.1007/s00701-014-2296-1>.
- [168] Eom KS. Single-stage reconstruction with titanium mesh for compound comminuted depressed skull fracture. *J Korean Neurosurg Soc* 2020;63:631–9. <https://doi.org/10.3340/jkns.2019.0181>.
- [169] Garst JR, Lara-Reyna J, Elliott W, Tsung AJ. Single-stage titanium mesh cranioplasty for the treatment of depressed skull fractures. *World Neurosurg* 2023;173:e62–5. <https://doi.org/10.1016/j.wneu.2023.01.103>.
- [170] Marbacher S, Andres RH, Fathi A-R, Fandino J. Primary reconstruction of open depressed skull fractures with titanium mesh. *J Craniofac Surg* 2008;19:490–5. <https://doi.org/10.1097/SCS.0b013e3181534ae8>.
- [171] Neville IS, Amorim RL, Paiva WS, Sanders FH, Teixeira MJ, de Andrade AF. Early surgery does not seem to be a pivotal criterion to improve prognosis in patients with frontal depressed skull fractures. *Biomed Res Int* 2014;2014:879286. <https://doi.org/10.1155/2014/879286>.
- [172] Al-Shami H, Alnemare AK, Mahfoz TB, Salah AM. Traumatic frontal sinus fractures management: experience from high-trauma centre. *Korean J Neurotrauma* 2021;17:15–24. <https://doi.org/10.13004/kjnt.2021.17.e3>.
- [173] Bell RB, Dierks EJ, Brar P, Potter JK, Potter BE. A protocol for the management of frontal sinus fractures emphasizing sinus preservation. *J Oral Maxillofac Surg* 2007;65:825–39. <https://doi.org/10.1016/j.joms.2006.05.058>.
- [174] Obayemi A, Losenegger T, Long S, Spielman D, Casiano MF, Reeve G, et al. Frontal sinus fractures: 10-year contemporary experience at a level 1 urban trauma center. *J Craniofac Surg* 2021;32:1376–80. <https://doi.org/10.1097/SCS.0000000000007426>.
- [175] Oslin K, Shikara M, Yoon J, Pope P, Bridgman K, Waghmare S, et al. Management of frontal sinus fractures at a level 1 trauma center: retrospective study and review of the literature. *Craniomaxillofac Trauma Reconstr* 2024;17:24–33. <https://doi.org/10.1177/19433875231155727>.
- [176] Afshari FT, Yakoub KM, Zisakis A, Thomas A, Ughratdar I, Sturman S, et al. Traumatic dural venous sinus thrombosis: a challenge in management of head injury patients. *J Clin Neurosci* 2018;57:169–73. <https://doi.org/10.1016/j.jocn.2018.08.047>.
- [177] Birk DM, Tobin MK, Moss HE, Feinstein E, Charbel FT, Alaraj A. Improvement in venous outflow following superior sagittal sinus decompression after a gunshot wound to the head: case report. *J Neurosurg* 2015;123:81–5. <https://doi.org/10.3171/2014.10.JNS141349>.
- [178] Forbes JA, Reig AS, Tomyecz LD, Tulipan N. Intracranial hypertension caused by a depressed skull fracture resulting in superior sagittal sinus thrombosis in a pediatric patient: treatment with ventriculoperitoneal shunt insertion. *J Neurosurg Pediatr* 2010;6:23–8. <https://doi.org/10.3171/2010.3.PEDS09441>.
- [179] Yokota H, Eguchi T, Nobayashi M, Nishiooka T, Nishimura F, Nikaido Y. Persistent intracranial hypertension caused by superior sagittal sinus stenosis following depressed skull fracture. Case report and review of the literature. *J Neurosurg* 2006;104:849–52. <https://doi.org/10.3171/jns.2006.104.5.849>.
- [180] Elshanawany A, Ragab M. Delayed neurological deterioration after depressed fracture over superior sagittal sinus: our experience with 13 patients. *J Neurol Surg A Cent Eur Neurosurg* 2020;81:324–9. <https://doi.org/10.1055/s-0039-1685189>.
- [181] Khalili H, Kashkooli NR, Niakan A, Asadi-Pooya AA. Risk factors for post-traumatic epilepsy. *Seizure* 2021;89:81–4. <https://doi.org/10.1016/j.seizure.2021.05.004>.
- [182] Oboth R, Kamabu LK, Lekuya HM, Bbosa GS, Sajatovic M, Katabira E, et al. Post-traumatic seizures and factors associated among adult patients with depressed skull fractures at Mulago National Referral hospital: cross-sectional study. *Epilepsy Behav* 2024;152:109693. <https://doi.org/10.1016/j.yebeh.2024.109693>.
- [183] Suliman A, Jayakumar N, Chaurasia B, Holliman D. Variations in the neurosurgical management of depressed skull fractures in adults: an international cross-sectional survey. *J Neurol Surg A Cent Eur Neurosurg* 2024;85:147–54. <https://doi.org/10.1055/a-1994-9330>.
- [184] Raveh J, Laedrach K, Vuillemin T, Zingg M. Management of combined frontonasal-orbital/skull base fractures and telecanthus in 355 cases. *Arch Otolaryngol Head Neck Surg* 1992;118:605–14. <https://doi.org/10.1001/archotol.1992.01880060053014>.
- [185] Scholsem M, Scholtes F, Collignon F, Robe P, Dubuisson A, Kaschten B, et al. Surgical management of anterior cranial base fractures with cerebrospinal fluid fistulae: a single-institution experience. *Neurosurgery* 2008;62:463–9. <https://doi.org/10.1227/01.neu.0000316014.97926.82>. discussion 469–471.
- [186] Sonig A, Thakur JD, Chittiboina P, Khan IS, Nanda A. Is posttraumatic cerebrospinal fluid fistula a predictor of posttraumatic meningitis? A US Nationwide Inpatient Sample database study. *Neurosurg Focus* 2012;32:E4. <https://doi.org/10.3171/2012.5.FOCUS1269>.
- [187] Oh J-W, Kim S-H, Whang K. Traumatic cerebrospinal fluid leak: diagnosis and management. *Korean J Neurotrauma* 2017;13:63–7. <https://doi.org/10.13004/kjnt.2017.13.2.63>.
- [188] Safavi A, Safavi AA, Jafari R. An empirical approach to the diagnosis and treatment of cerebrospinal fluid rhinorrhea: an optimised method for developing countries. *Malays J Med Sci* 2014;21:37–43.
- [189] Shorter CD, Connor DE, Thakur JD, Gardner G, Nanda A, Guthikonda B. Repair of middle fossa cerebrospinal fluid leaks using a novel combination of materials: technical note. *Neurosurg Focus* 2012;32:E8. <https://doi.org/10.3171/2012.4.FOCUS1258>.
- [190] Banks CA, Palmer JN, Chiu AG, O'Malley BW, Woodworth BA, Kennedy DW. Endoscopic closure of CSF rhinorrhea: 193 cases over 21 years. *Otolaryngol Head Neck Surg* 2009;140:826–33. <https://doi.org/10.1016/j.otohns.2008.12.060>.
- [191] Themes UFO. Traumatic Cerebrospinal Fluid Fistulas. Neupsy Key 2016. <https://neupsykey.com/traumatic-cerebrospinal-fluid-fistulas/> (accessed August 20, 2024).
- [192] Lindstrom DR, Toohill RJ, Loehrl TA, Smith TL. Management of cerebrospinal fluid rhinorrhea: the Medical College of Wisconsin experience. *Laryngoscope* 2004;114:969–74. <https://doi.org/10.1097/00005537-200406000-00003>.
- [193] Yilmazlar S, Arslan E, Kocaeli H, Dogan S, Aksoy K, Korfali E, et al. Cerebrospinal fluid leakage complicating skull base fractures: analysis of 81 cases. *Neurosurg Rev* 2006;29:64–71. <https://doi.org/10.1007/s10143-005-0396-3>.
- [194] Sherif C, Di Ieva A, Gibson D, Pakrah-Bodingbauer B, Widhalm G, Krusche-Mandl I, et al. A management algorithm for cerebrospinal fluid leak associated with anterior skull base fractures: detailed clinical and radiological follow-up. *Neurosurg Rev* 2012;35:227–37. <https://doi.org/10.1007/s10143-011-0352-3>. discussion 237–238.
- [195] Prosser JD, Vender JR, Solares CA. Traumatic cerebrospinal fluid leaks. *Otolaryngol Clin North Am* 2011;44:857–73. <https://doi.org/10.1016/j.otc.2011.06.007>. vii.
- [196] Rocchi G, Caroli E, Belli E, Salvati M, Cimatti M, Delfini R. Severe craniofacial fractures with frontobasal involvement and cerebrospinal fluid fistula: indications for surgical repair. *Surg Neurol* 2005;63:559–63. <https://doi.org/10.1016/j.surneu.2004.07.047>. discussion 563–564.
- [197] Eljamel MS, Foy PM. Post-traumatic CSF fistulae, the case for surgical repair. *Br J Neurosurg* 1990;4:479–83. <https://doi.org/10.1019/02688699008993796>.
- [198] Friedman JA, Ebersold MJ, Quast LM. Post-traumatic cerebrospinal fluid leakage. *World J Surg* 2001;25:1062–6. <https://doi.org/10.1007/s00268-001-0059-7>.
- [199] Fishman G, Fliss DM, Benjamin S, Margalit N, Gil Z, Derowe A, et al. Multidisciplinary surgical approach for cerebrospinal fluid leak in children with complex head trauma. *Childs Nerv Syst* 2009;25:915–23. <https://doi.org/10.1007/s00381-009-0887-6>.
- [200] Schoentgen C, Henaux PL, Godey B, Jegoux F. Management of post-traumatic cerebrospinal fluid (CSF) leak of anterior skull base: 10 years experience. *Acta Otolaryngol* 2013;133:944–50. <https://doi.org/10.3109/00016489.2013.793821>.
- [201] Thapa AJ, Lei B-X, Zheng M-G, Li Z-J, Liu Z-H, Deng Y-F. The surgical treatment of posttraumatic skull base defects with cerebrospinal fluid leak. *J Neurol Surg B Skull Base* 2018;79:205–16. <https://doi.org/10.1055/s-0037-1606555>.
- [202] Fliss DM, Zucker G, Cohen JT, Gatot A. The subcranial approach for the treatment of cerebrospinal fluid rhinorrhea: a report of 10 cases. *J Oral Maxillofac Surg* 2001;59:1171–5. <https://doi.org/10.1053/joms.2001.26720>.
- [203] Talamonti G, Fontana R, Villa F, D'Aliberti G, Arena O, Bizzozero L, et al. "High risk" anterior basal skull fractures. Surgical treatment of 64 consecutive cases. *J Neurosurg Sci* 1995;39:191–7.
- [204] Archer JB, Sun H, Bonney PA, Zhao YD, Hiebert JC, Sanclement JA, et al. Extensive traumatic anterior skull base fractures with cerebrospinal fluid leak: classification and repair techniques using combined vascularized tissue flaps. *J Neurosurg* 2016;124:647–56. <https://doi.org/10.3171/2015.4.JNS1528>.
- [205] Bell RB, Dierks EJ, Homer L, Potter BE. Management of cerebrospinal fluid leak associated with craniomaxillofacial trauma. *J Oral Maxillofac Surg* 2004;62:676–84. <https://doi.org/10.1016/j.joms.2003.08.032>.
- [206] Dalgic A, Okay HO, Gezici AR, Daglioglu E, Akdag R, Ergunor MF. An effective and less invasive treatment of post-traumatic cerebrospinal fluid fistula: closed lumbar drainage system. *Minim Invasive Neurosurg* 2008;51:154–7. <https://doi.org/10.1055/s-2008-1042437>.

- [207] Albu S, Florian IS, Bolboaca SD. The benefit of early lumbar drain insertion in reducing the length of CSF leak in traumatic rhinorrhea. *Clin Neurol Neurosurg* 2016;142:43–7. <https://doi.org/10.1016/j.clineuro.2016.01.019>.
- [208] Khan R, Sajjad M, Khan AA, Ahmad B, Mushtaq M, et al. Comparison Of Lumbar Drain Insertion And Conservative Management In The Treatment Of Traumatic CSF Rhinorrhea. *J Ayub Med Coll Abbottabad* 2019;31:441–4.
- [209] Recommendations SFAR SPILF 2023 - Antibiotrophylaxie en chirurgie et médecine interventionnelle. <https://sfar.org/antibiotrophylaxie-en-chirurgie-et-medecine-interventionnelle/>. Accès le 14/12/2023. n.d.
- [210] Ter Horst L, Brouwer MC, van der Ende A, van de Beek D. Community-acquired bacterial meningitis in adults with cerebrospinal fluid leakage. *Clin Infect Dis* 2020;70:2256–61. <https://doi.org/10.1093/cid/ciz649>.
- [211] Eljazzar R, Loewenstern J, Dai JB, Shrivastava RK, Illoreta AM. Detection of cerebrospinal fluid leaks: is there a radiologic standard of care? A systematic review. *World Neurosurgery* 2019;127:307–15. <https://doi.org/10.1016/j.wneu.2019.01.299>.
- [212] Xie M, Zhou K, Kachra S, McHugh T, Sommer DD. Diagnosis and localization of cerebrospinal fluid rhinorrhea: a systematic review. *Am J Rhinol Allergy* 2022;36:397–406. <https://doi.org/10.1177/19458924211060918>.
- [213] Aarabi B, Tofighi B, Kufera JA, Hadley J, Ahn ES, Cooper C, et al. Predictors of outcome in civilian gunshot wounds to the head. *J Neurosurg* 2014;120:1138–46. <https://doi.org/10.3171/2014.1.JNS131869>.
- [214] Alvis-Miranda H.R., Rubiano A.M., Agrawal A., Rojas A., Rafael L., Satyarthee G. D., et al. Craniocerebral Gunshot Injuries: A Review of the Current Literature n.d.
- [215] Kim T-W, Lee J-K, Moon K-S, Kwak H-J, Joo S-P, Kim J-H, et al. Penetrating gunshot injuries to the brain. *J Trauma* 2007;62:1446–51. <https://doi.org/10.1097/01.ta.0000222909.31666.dy>.
- [216] Mansour A, Powla PP, Alvarado-Dyer R, Fakhri F, Das P, Horowitz P, et al. Comparative analysis of clinical severity and outcomes in penetrating versus blunt traumatic brain injury propensity matched cohorts. *Neurotrauma Rep* 2024;5:348–58. <https://doi.org/10.1089/neur.2024.0009>.
- [217] Robinson LA, Turco LM, Robinson B, Corsa JG, Mount M, Hamrick AV, et al. Outcomes in patients with gunshot wounds to the brain. *Trauma Surg Acute Care Open* 2019;4:e000351. <https://doi.org/10.1136/tsaco-2019-000351>.
- [218] Rosenfeld JV, Bell RS, Armonda R. Current concepts in penetrating and blast injury to the central nervous system. *World J Surg* 2015;39:1352–62. <https://doi.org/10.1007/s00268-014-2874-7>.
- [219] Vakili MT, Singh AK. A review of penetrating brain trauma: epidemiology, pathophysiology, imaging assessment, complications, and treatment. *Emerg Radiol* 2017;24:301–9. <https://doi.org/10.1007/s10140-016-1477-z>.
- [220] Dagain A, Aoun O, Bordes J, Roqueplo C, Joubert C, Esnault P, et al. Management of war-related ballistic craniocerebral injuries in a french role 3 hospital during the Afghan campaign. *World Neurosurg* 2017;102:6–12. <https://doi.org/10.1016/j.wneu.2017.02.097>.
- [221] Dagain A, Aoun O, Sellier A, Desse N, Joubert C, Beucler N, et al. Acute neurosurgical management of traumatic brain injury and spinal cord injury in French armed forces during deployment. *Neurosurg Focus* 2018;45:E9. <https://doi.org/10.3171/2018.9.FOCUS18368>.
- [222] Rosenfeld JV. Gunshot injury to the head and spine. *J Clin Neurosci* 2002;9:9–16. <https://doi.org/10.1054/jocn.2001.0949>.
- [223] Gressot LV, Chamoun RB, Patel AJ, Valadka AB, Suki D, Robertson CS, et al. Predictors of outcome in civilians with gunshot wounds to the head upon presentation: clinical article. *JNS* 2014;121:645–52. <https://doi.org/10.3171/2014.5.JNS131872>.
- [224] Maragkos GA, Papavassiliou E, Stippler M, Filippidis AS. Civilian gunshot wounds to the head: prognostic factors affecting mortality: meta-analysis of 1774 patients. *J Neurotrauma* 2018;35:2605–14. <https://doi.org/10.1089/neu.2018.5682>.
- [225] Kong V, Odendaal J, Sartorius B, Clarke D, Bruce J, Laing G, et al. Developing a simplified clinical prediction score for mortality in patients with cerebral gunshot wounds: the Maritzburg Score. *Annals* 2018;100:97–100. <https://doi.org/10.1308/rcsann.2017.0141>.
- [226] Muehlschlegel S, Ayturk D, Ahlawat A, Izzy S, Scalea TM, Stein DM, et al. Predicting survival after acute civilian penetrating brain injuries: the SPIN score. *Neurology* 2016;87:2244–53. <https://doi.org/10.1212/WNL.0000000000003355>.
- [227] Levy ML. Outcome prediction following penetrating craniocerebral injury in a civilian population: aggressive surgical management in patients with admission Glasgow Coma Scale scores of 6 to 15. *Neurosurg Focus* 2000;8:e2. <https://doi.org/10.3171/foc.2000.8.1.153>.
- [228] Petridis AK, Doukas A, Barth H, Mehdorn M. Outcome of craniocerebral gunshot injuries in the civilian population. Prognostic Factors and Treatment Options. *Cen Eur Neurosurg* 2011;72:5–14. <https://doi.org/10.1055/s-0029-1241850>.
- [229] Hubschmann O, Shapiro K, Baden M, Shulman K. Craniocerebral gunshot injuries in civilian practice—prognostic criteria and surgical management: experience with 82 cases. *J Trauma* 1979;19:6–12.
- [230] Helling TS, McNABNEY WK, Whittaker CK, Schultz CC, Watkins M. The role of early surgical intervention in civilian gunshot wounds to the head. *J Trauma* 1992;32:398–400. <https://doi.org/10.1097/00005373-199203000-00019>.
- [231] Surgical management of penetrating brain injury. *J Trauma* 2001;51:S16–25.
- [232] Esposito DP, Walker JB. Contemporary management of penetrating brain injury. *Neurosurg Q* 2009;19:249–54. <https://doi.org/10.1097/WNQ.0b013e3181b1d153>.
- [233] Bizhan A, Musa T, Ehsanali A, Ahmad K, John CF. Central nervous system infections after military missile head wounds. *Neurosurgery* 1998;42:500–9. <https://doi.org/10.1097/00006123-199803000-00014>.
- [234] Gönül E, Baysefer A, Kahraman S, Çklatekerlioğlu Ö, Gezen F, Yayla O, et al. Causes of infections and management results in penetrating craniocerebral injuries. *Neurosurg Rev* 1997;20:177–81. <https://doi.org/10.1007/BF01105561>.
- [235] Chaudhri KA, Choudhury AR, Al Moutaery KR, Cybulski GR. Penetrating craniocerebral shrapnel injuries during “Operation Desert Storm”: early results of a conservative surgical treatment. *Acta Neurochir* 1994;126:120–3. <https://doi.org/10.1007/BF01476420>.
- [236] Salazar AM, Jabbari B, Vance SC, Grafman J, Amin D, Dillon JD. Epilepsy after penetrating head injury. I. Clinical correlates: A report of the Vietnam Head Injury Study. *Neurology* 1985;35:1406. <https://doi.org/10.1212/WNL.35.10.1406>.
- [237] Hammon WM. Analysis of 2187 consecutive penetrating wounds of the brain from Vietnam. *J Neurosurg* 1971;34:127–31. <https://doi.org/10.3171/jns.1971.34.2part1.0127>.
- [238] Manet R, Joubert C, Balanca B, Taverna X-J, Monneuse O, David J-S, et al. Neuro damage control: current concept and civilian applications. *Neurochirurgie* 2023;69:101505. <https://doi.org/10.1016/j.neuchi.2023.101505>.
- [239] Gutiérrez-González R, Boto GR, Rivero-Garvía M, Pérez-Zamarrón Á, Gómez G. Penetrating brain injury by drill bit. *Clin Neurol Neurosurg* 2008;110:207–10. <https://doi.org/10.1016/j.clineuro.2007.09.014>.
- [240] Rish BL, Dillon JD, Caveness WF, Mohr JP, Kistler JP, Weiss GH. Evolution of craniotomy as a debridement technique for penetrating craniocerebral injuries. *J Neurosurg* 1980;53:772–5. <https://doi.org/10.3171/jns.1980.53.6.0772>.
- [241] Bell RS, Vo AH, Roberts R, Wanebo J, Armonda RA. Wartime traumatic aneurysms: acute presentation, diagnosis, and multimodal treatment of 64 craniocervical arterial injuries. *Neurosurgery* 2010;66:66–79. <https://doi.org/10.1227/01.NEU.0000361285.50218.A8>. discussion 79.
- [242] Krueger EM, Moll J, Kumar R, Lu VM, Benveniste R, Cordeiro JG, et al. Simple wound closure for civilian cranial gunshot wounds: a systematic literature review. *Cureus* 2022. <https://doi.org/10.7759/cureus.25187>.
- [243] Souza RBD, Todeschini AB, Veiga JCE, Saade N, Aguiar GBD. Traumatismo cranioencefálico por projétil de arma de fogo: experiência de 16 anos do serviço de neurocirurgia da Santa Casa de São Paulo. *Rev Col Bras Cir* 2013;40:300–4. <https://doi.org/10.1590/S0100-69912013000400008>.
- [244] Loggini A, Vasenina VI, Mansour A, Das P, Horowitz PM, Goldenberg FD, et al. Management of civilians with penetrating brain injury: a systematic review. *J Crit Care* 2020;56:159–66. <https://doi.org/10.1016/j.jcrr.2019.12.026>.
- [245] Walker WC, Ketchum JS, Marwitz JH, Kolakowsky-Hayner SA, McClish DK, Bushnik T. Global outcome and late seizures after penetrating versus closed traumatic brain injury: a NIDRR TBI model systems study. *J Head Trauma Rehabil* 2015;30:231–40. <https://doi.org/10.1097/HTR.0000000000000127>.
- [246] Karras CL, Texakalidis P, Nie JZ, Tran HM, Dahdaleh NS, Bovis GK, et al. Outcomes following penetrating brain injuries in military settings: a systematic review and meta-analysis. *World Neurosurg* 2022;166:39–48. <https://doi.org/10.1016/j.wneu.2022.07.062>.
- [247] Picetti E, Rossi S, Abu-Zidan FM, Ansaloni L, Armonda R, Baiocchi GL, et al. WSES consensus conference guidelines: monitoring and management of severe adult traumatic brain injury patients with polytrauma in the first 24 hours. *World J Emerg Surg* 2019;14:53. <https://doi.org/10.1186/s13017-019-0270-1>.
- [248] Antiseizure prophylaxis for penetrating brain injury. *J Trauma* 2001;51:S41–43.
- [249] Mansour A, Loggini A, El Ammar F, Ginat D, Awad IA, Lazaridis C, et al. Cerebrovascular complications in early survivors of civilian penetrating brain injury. *Neurocrit Care* 2021;34:918–26. <https://doi.org/10.1007/s12028-020-01106-y>.
- [250] Szmuda T, Sloniewski P, Waszak PM, Springer J, Szmuda M. Towards a new treatment paradigm for ruptured blood blister-like aneurysms of the internal carotid artery? A rapid systematic review. *J Neurointerv Surg* 2016;8:488–94. <https://doi.org/10.1136/neurintsurg-2015-011665>.
- [251] Meyer RM, Grandhi R, Lim DH, Salah WK, McAvoy M, Abecassis ZA, et al. A comparison of computed tomography angiography and digital subtraction angiography for the diagnosis of penetrating cerebrovascular injury: a prospective multicenter study. *J Neurosurg* 2024;141:306–9. <https://doi.org/10.3171/2023.12.JNS232070>.
- [252] Ares WJ, Jankowitz BT, Tonetti DA, Gross BA, Grandhi R. A comparison of digital subtraction angiography and computed tomography angiography for the diagnosis of penetrating cerebrovascular injury. *Neurosurg Focus* 2019;47:E16. <https://doi.org/10.3171/2019.8.FOCUS19495>.
- [253] Amirjamshidi A, Rahmat H, Abbassioun K. Traumatic aneurysms and arteriovenous fistulas of intracranial vessels associated with penetrating head injuries occurring during war: principles and pitfalls in diagnosis and management. A survey of 31 cases and review of the literature. *J Neurosurg* 1996;84:769–80. <https://doi.org/10.3171/jns.1996.84.5.0769>.
- [254] Lefevre E, Fawaz R, Premat K, Lenck S, Shoter E, Degos V, et al. Delayed traumatic intracranial aneurysms: literature review and case series. *Neurosurg Rev* 2024;47(1):355. <https://doi.org/10.1007/s10143-024-02586-8>. PMID: 39060452.
- [255] Bell RS, Ecker RD, Severson 3rd MA, Wanebo JE, Crandall B, Armonda RA. The evolution of the treatment of traumatic cerebrovascular injury during wartime. *Neurosurg Focus* 2010;28(5):E5. <https://doi.org/10.3171/2010.2.FOCUS1025>. PMID: 20568945.
- [256] Stone JL, Lang RG, Sugar O, Moody RA. Traumatic subdural hygroma. *Neurosurgery* 1981;8:542–50. <https://doi.org/10.1227/00006123-198105000-00005>.
- [257] Naffziger HC. Subdural fluid accumulations following head injury. *J Am Med Assoc* 1924;82:1751–2. <https://doi.org/10.1001/jama.1924.02650480007004>.
- [258] García Romero JC, Ortega Martínez R, Zabalo San Juan G, de Frutos Marcos D, Zazpe Cenoz I. Subdural hygroma secondary to rupture of an intracranial

- arachnoid cyst: description of 2 cases and review of the literature. *Neurocirugia (Astur: Engl Ed)* 2018;29:260–4. <https://doi.org/10.1016/j.neucir.2018.02.003>.
- [259] Gelabert-González M, Fernández-Villa J, Cutrín-Prieto J, García Allut A, Martínez-Rumbo R. Arachnoid cyst rupture with subdural hygroma: report of three cases and literature review. *Childs Nerv Syst* 2002;18:609–13. <https://doi.org/10.1007/s00381-002-0651-7>.
- [260] Gupta R, Vaishya S, Mehta VS. Arachnoid cyst presenting as subdural hygroma. *J Clin Neurosci* 2004;11:317–8. <https://doi.org/10.1016/j.jocn.2003.03.003>.
- [261] Tamburrini G, Caldarelli M, Massimi L, Santini P, Di Rocco C. Subdural hygroma: an unwanted result of Sylvian arachnoid cyst marsupialization. *Childs Nerv Syst* 2003;19:159–65. <https://doi.org/10.1007/s00381-003-0724-2>.
- [262] Lee KS. The pathogenesis and clinical significance of traumatic subdural hygroma. *Brain Inj* 1998;12:595–603. <https://doi.org/10.1080/026990598122359>.
- [263] Zanini MA, de Lima Resende LA, de Souza Faleiros AT, Gabarra RC. Traumatic subdural hygromas: proposed pathogenesis based classification. *J Trauma* 2008;64:705–13. <https://doi.org/10.1097/TA.0b013e3180485cfc>.
- [264] Gergely L, Manet R, Kolias A, Czosnyka M, Lalou A, Smielewski P, et al. External hydrocephalus after traumatic brain injury: retrospective study of 102 patients. *Acta Neurochir Suppl* 2021;131:35–8. https://doi.org/10.1007/978-3-030-59436-7_8.
- [265] Manet R, Payen J-F, Guerin R, Martinez O, Hautefeuille S, Francony G, et al. Using external lumbar CSF drainage to treat communicating external hydrocephalus in adult patients after acute traumatic or non-traumatic brain injury. *Acta Neurochir (Wien)* 2017;159:2003–9. <https://doi.org/10.1007/s00701-017-3290-1>.
- [266] Huh P-W, Yoo D-S, Cho K-S, Park K-K, Kang S-G, Park Y-S, et al. Diagnostic method for differentiating external hydrocephalus from simple subdural hygroma. *J Neurosurg* 2006;105:65–70. <https://doi.org/10.3171/jns.2006.105.1.65>.
- [267] Kaen A, Jimenez-Roldan L, Alday R, Gomez PA, Lagares A, Alén JF, et al. Inter-hemispheric hygroma after decompressive craniectomy: does it predict post-traumatic hydrocephalus? *J Neurosurg* 2010;113:1287–93. <https://doi.org/10.3171/2010.4.JNS10132>.
- [268] Kim JH, Ahn JH, Oh JK, Song JH, Park SW, Chang IB. Factors associated with the development and outcome of hydrocephalus after decompressive craniectomy for traumatic brain injury. *Neurosurg Rev* 2021;44:471–8. <https://doi.org/10.1007/s10143-019-01179-0>.
- [269] Paredes I, Cicuendez M, Delgado MA, Martínez-Pérez R, Munarriz PM, Lagares A. Normal pressure subdural hygroma with mass effect as a complication of decompressive craniectomy. *Surg Neurol Int* 2011;2:88. <https://doi.org/10.4103/2152-7806.82370>.
- [270] Shen X, Han Y, Li H, Dong Y, Yang D, Xu W, et al. The risk factors associated with traumatic subdural effusion for patients with traumatic brain injury who did not undergo decompressive craniectomy. *Acta Neurol Belg* 2023;123:957–61. <https://doi.org/10.1007/s13760-022-02099-5>.
- [271] Vedantam A, Yamal J-M, Hwang H, Robertson CS, Gopinath SP. Factors associated with shunt-dependent hydrocephalus after decompressive craniectomy for traumatic brain injury. *J Neurosurg* 2018;128:1547–52. <https://doi.org/10.3171/2017.1.JNS162721>.
- [272] Murad A, Ghostine S, Colohan ART. Controlled lumbar drainage in medically refractory increased intracranial pressure. A safe and effective treatment. *Acta Neurochir Suppl* 2008;102:89–91. https://doi.org/10.1007/978-3-211-85578-2_18.
- [273] Tuettenberg J, Czabanka M, Horn P, Woitzik J, Barth M, Thomé C, et al. Clinical evaluation of the safety and efficacy of lumbar cerebrospinal fluid drainage for the treatment of refractory increased intracranial pressure. *J Neurosurg* 2009;110:1200–8. <https://doi.org/10.3171/2008.10.JNS08293>.
- [274] Soltész R, Dömötör H, Varga ÁL, Marada M, Baracska E, Radványi S, et al. Controlled lumbar cerebrospinal fluid drainage effectively decreases the need for second and third tier interventions for intracranial hypertension in severe traumatic brain injury patients. *Injury* 2024;55(Suppl 3):111337. <https://doi.org/10.1016/j.injury.2024.111337>.
- [275] Bauer M, Sohm F, Thomé C, Ortler M. Refractory intracranial hypertension in traumatic brain injury: proposal for a novel score to assess the safety of lumbar cerebrospinal fluid drainage. *Surg Neurol Int* 2017;8:265. <https://doi.org/10.4103/sni.sni.98.17>.
- [276] Grille P, Biestro A, Rekaté HL. Intracranial hypertension with patent basal cisterns: controlled lumbar drainage as a therapeutic option. Selected case series. *Neurocrit Care* 2024;40:1070–82. <https://doi.org/10.1007/s12028-023-01878-z>.
- [277] Llompart-Pou JA, Abadal JM, Pérez-Bárcena J, Molina M, Brell M, Ibáñez J, et al. Long-term follow-up of patients with post-traumatic refractory high intracranial pressure treated with lumbar drainage. *Anaesth Intensive Care* 2011;39:79–83. <https://doi.org/10.1177/0310057X1103900113>.
- [278] Stevens AR, Soon WC, Chowdhury YA, Toman E, Yim S, Veenith T, et al. External lumbar drainage for refractory intracranial hypertension in traumatic brain injury: a systematic review. *Cureus* 2022;14:e30033. <https://doi.org/10.7759/cureus.30033>.
- [279] Levy DI, Rekaté HL, Cherny WB, Manwaring K, Moss SD, Baldwin HZ. Controlled lumbar drainage in pediatric head injury. *J Neurosurg* 1995;83:453–60. <https://doi.org/10.3171/jns.1995.83.3.0453>.
- [280] Willemse RB, Egeler-Peerdeman SM. External lumbar drainage in uncontrollable intracranial pressure in adults with severe head injury: a report of 7 cases. *Acta Neurochir Suppl* 1998;71:37–9. https://doi.org/10.1007/978-3-7091-6475-4_11.
- [281] Münch EC, Bauhuf C, Horn P, Roth HR, Schmiedek P, Vajkoczy P. Therapy of malignant intracranial hypertension by controlled lumbar cerebrospinal fluid drainage. *Crit Care Med* 2001;29:976–81. <https://doi.org/10.1097/00003246-200105000-00016>.
- [282] Manet R, Schmidt EA, Vassal F, Chariot D, Gergely L. CSF lumbar drainage: a safe surgical option in refractory intracranial hypertension associated with acute posttraumatic external hydrocephalus. *Acta Neurochir Suppl* 2016;122:55–9. https://doi.org/10.1007/978-3-319-22533-3_11.
- [283] Abadal-Centellas JM, Llompart-Pou JA, Homar-Ramírez J, Pérez-Bárcena J, Rosselló-Ferrer A, Ibáñez-Juvé J. Neurologic outcome of posttraumatic refractory intracranial hypertension treated with external lumbar drainage. *J Trauma* 2007;62:282–6. <https://doi.org/10.1097/01.ta.0000199422.01949.78>. discussion 286.
- [284] Tömösvári A, Mencser Z, Futó J, Hortobágyi A, Bodosi M, Barzó P. [Preliminary experience with controlled lumbar drainage in medically refractory intracranial hypertension]. *Orv Hetil* 2005;146:159–64.
- [285] Iranmehr A, Rashidbeygi M, Namvar M, Khadivi A, Hoseynzadeh E, Payinmahalli A. Neonatal acute epidural hematoma: a case report and literature review. *Korean J Neurotrauma* 2020;16:262. <https://doi.org/10.13004/kjnt.2020.16.e22>.
- [286] Paiva WS, Andrade AF de, Mathias Júnior L, Guirado VM de P, Amorim RL, Magrini NN, et al. Management of supratentorial epidural hematoma in children: report on 49 patients. *Arq Neuropsiquiatr* 2010;68:888–92. <https://doi.org/10.1590/s0004-282x2010000600011>.
- [287] Baş NS, Karacan M, Doruk E, Karagoz Guzey F. Management of traumatic epidural hematoma in infants younger than one year: 50 cases - single center experience. *Pediatr Neurosurg* 2021;56:213–20. <https://doi.org/10.1159/000514810>.
- [288] Irie F, Le Brocq R, Kenardy J, Bellamy N, Tetsworth K, Pollard C. Epidemiology of traumatic epidural hematoma in young age. *J Trauma* 2011;71:847–53. <https://doi.org/10.1097/TA.0b013e3182032c9a>.
- [289] Jamous MA, Abdel Aziz H, Al Kaisy F, Eloqayli H, Azab M, Al-Jarrah M. Conservative management of acute epidural hematoma in a pediatric age group. *Pediatr Neurosurg* 2009;45:181–4. <https://doi.org/10.1159/000218200>.
- [290] Vachharajani A, Mathur A. Ultrasound-guided needle aspiration of cranial epidural hematoma in a neonate: treating a rare complication of vacuum extraction. *Am J Perinatol* 2002;19:401–4. <https://doi.org/10.1055/s-2002-36837>.
- [291] Hamlat A, Heckly A, Adn M, Poulain P. Pathophysiology of intracranial epidural haematoma following birth. *Med Hypotheses* 2006;66:371–4. <https://doi.org/10.1016/j.mehy.2005.08.045>.
- [292] Bejjani GK, Donahue DJ, Rusin J, Broemeling LD. Radiological and clinical criteria for the management of epidural hematomas in children. *Pediatr Neurosurg* 1996;25:302–8. <https://doi.org/10.1159/000121144>.
- [293] Smets KJ, Vanhauwaert D. Treatment of cranial epidural hematoma in a neonate by needle aspiration of a communicating cephalhematoma. *Eur J Pediatr* 2010;169:617–9. <https://doi.org/10.1007/s00431-009-1071-4>.
- [294] Govaert P. Epidural haematoma (cephalhematoma internum, internal sub-periosteal bleeding). *Cranial Haemorrhage in the Term Newborn Infant. Clinics in Developmental Medicine*. Cambridge University Press; 1994. p. 34–40.
- [295] Heyman R, Heckly A, Magagi J, Pladys P, Hamlat A. Intracranial epidural hematoma in newborn infants: clinical study of 15 cases. *Neurosurgery* 2005;57:924–9. <https://doi.org/10.1227/01.NEU.0000180026.73246.bf>.
- [296] Vinchon M, Noulé N, Soto-Ares G, Dhellemmes P. Subduroperitoneal drainage for subdural hematomas in infants: results in 244 cases. *J Neurosurg* 2001;95:249–55. <https://doi.org/10.3171/jns.2001.95.2.0249>.
- [297] Melo JRT, Di Rocco F, Bourgeois M, Puget S, Blauwblomme T, Sainte-Rose C, et al. Surgical options for treatment of traumatic subdural hematomas in children younger than 2 years of age: clinical article. *PED* 2014;13:456–61. <https://doi.org/10.3171/2014.1.PEDS13393>.
- [298] Vinchon M, Defoort-Dhellemmes S, Desurmont M, Dhellemmes P. Accidental and nonaccidental head injuries in infants: a prospective study. *J Neurosurg* 2005;102:380–4. <https://doi.org/10.3171/2005.102.4.0380>.
- [299] Kochanek PM, Tasker RC, Bell MJ, Adelson PD, Carney N, Vavilala MS, et al. Management of pediatric severe traumatic brain injury: 2019 consensus and guidelines-based algorithm for first and second tier therapies. *Pediatr Crit Care Med* 2019;20:269–79. <https://doi.org/10.1097/PCC.0000000000001737>.
- [300] Karibe H, Kameyama M, Hayashi T, Narisawa A, Tominaga T. Acute subdural hematoma in infants with abusive head trauma: a literature review. *Neurol Med Chir (Tokyo)* 2016;56:264–73. <https://doi.org/10.2176/nmc.ra.2015-0308>.
- [301] Elsawaf Y, Anetsberger S, Luzzi S, Elbabaa SK. Early decompressive craniectomy as management for severe traumatic brain injury in the pediatric population: a comprehensive literature review. *World Neurosurg* 2020;138:9–18. <https://doi.org/10.1016/j.wneu.2020.02.065>.
- [302] Taylor A, Butt W, Rosenfeld J, Shann F, Ditchfield M, Lewis E, et al. A randomized trial of very early decompressive craniectomy in children with traumatic brain injury and sustained intracranial hypertension. *Child's Nervous System* 2001;17:154–62. <https://doi.org/10.1007/s003810000410>.
- [303] Jagannathan J, Okonkwo DO, Dumont AS, Ahmed H, Bahari A, Prevedello DM, et al. Outcome following decompressive craniectomy in children with severe traumatic brain injury: a 10-year single-center experience with long-term follow up. *J Neurosurg* 2007;106:268–75. <https://doi.org/10.3171/ped.2007.106.4.268>.
- [304] Altieri R, Grasso E, Cammarata G, Garozzo M, Marchese G, Certo F, et al. Decision-making challenge of ping-pong fractures in children: case exemplification and systematic review of literature. *World Neurosurg* 2022;165:69–80. <https://doi.org/10.1016/j.wneu.2022.05.130>.
- [305] Ahmed SD, Allhusen VD, Muhonen MG, Magge SN. Pediatric ping-pong skull fractures treated with vacuum-assisted elevation. *Childs Nerv Syst* 2024;40:1833–8. <https://doi.org/10.1007/s00381-024-06307-w>.

Romain Manet^{a,b,*}, Hugues de Courson^{c,d}, Cyrille Capel^{e,f},
 Christophe Joubert^{b,g}, Nathalie Chivoret^h, Matthieu Faillot^{i,j},
 Baptiste Balanca^{k,l}, Alexandre Bani-Sadr^{m,n}, Mickael Cardinale^{g,o},
 Andres Coca^p, François Cotton^{n,q}, Pierre Esnault^o, Clémentine Gallet^a,
 Sébastien Gazzola^r, Stéphane Goutagny^{h,s}, Vincent Jecko^{t,u}, Marion le
 Marechal^{v,w}, Jacques Luauté^{x,y}, Guillaume Mortamet^{w,z}, Jean-
 Denis Moyer^A, Hervé Quintard^B, Alice Rolland^C, Édouard Samarut^{D,E},
 Stéphanie Sigaut^{F,G}, Eric Verin^{H,I}, Mathieu Vinchon^J, Philippe Decq^{I,K},
 Jean-François Payen^w, Arnaud Dagain^{b,g}

^a Service de Neurochirurgie Crânienne, Hôpital Neurologique Pierre
 Wertheimer, Hospices Civils de Lyon, Bron, France

^b Service de Neurochirurgie, Hôpital National d'Instruction des Armées
 Saint-Anne, Toulon, France

^c Département d'Anesthésie-Réanimation, Hôpital Pellegrin, CHU de
 Bordeaux, Bordeaux, France

^d Université de Bordeaux, INSERM UMR 1219, Bordeaux, France

^e Service de Neurochirurgie, CHU Amiens-Picardie, Amiens, France

^f CHIMERE UR UPJV 7516, Université de Picardie Jules Verne, Amiens,
 France

^g Ecole du Val de Grâce, Académie de Santé des Armées, Paris, France

^h Service de Neurochirurgie, CHU de Nice, Nice, France

ⁱ Service de Neurochirurgie, Hôpital Beaujon, Assistance Publique Hôpitaux
 de Paris, Clichy, France

^j Université Paris-Cité, Laboratoire CEA/Service Hospitalier FrédéricJoliot/
 BioMaps, Paris, France

^k Service d'Anesthésie Réanimation Neurologique, Hôpital Neurologique
 Pierre Wertheimer, Hospices Civils de Lyon, Bron, France

^l Centre de Recherche en Neurosciences, INSERM U1028/CNRS, UMR
 5292, Université de Lyon 1, Lyon, France

^m Service de Neuroradiologie, Hôpital Neurologique Pierre Wertheimer,
 Hospices Civils de Lyon, Bron, France

ⁿ CREATIS-LRMN, CNRS/UMR/5220-INSERM U630, Université Lyon 1,
 Lyon, France

^o Service de Réanimation, Hôpital National d'Instruction des Armées Sainte
 Anne, Toulon, France

^p Service de Neurochirurgie, Hôpital Hautepierre, CHU Strasbourg,
 Strasbourg, France

^q Service de Radiologie, Centre Hospitalier Lyon Sud, Hospices Civils de
 Lyon, Lyon, France

^r Service de Neurologie, Hôpital National d'Instruction des Armées Sainte
 Anne, Toulon, France

^s Université Paris Cité, Inserm UMRS1144, Paris, France

^t Service de Neurochirurgie A, Hôpital Pellegrin, CHU Bordeaux, Bordeaux,
 France

^u Université de Bordeaux, CNRS UMR, INCIA, Bordeaux, France

^v Service de Maladies Infectieuses, CHU Grenoble Alpes, Grenoble, France

^w Université Grenoble-Alpes, Grenoble, France

^x Service de Médecine Physique et Réadaptation, Hôpital Henry Gabrielle,
 Hospices Civils de Lyon, Saint-Genis-Laval, France

^y Centre de Recherche en Neurosciences de Lyon, U1028, UMR5229, équipe
 Trajectoires, Université Lyon 1, Lyon, France

^z Service de Soins Critiques Pédiatriques, CHU de Grenoble, Grenoble,
 France

^A Département d'Anesthésie-Réanimation, CHU de Caen Normandie, Caen,
 France

^B Département d'Anesthésiologie, de Pharmacologie Clinique, de Soins
 Intensifs et de Médecine d'Urgence, Hôpitaux Universitaires de Genève,
 Genève, Switzerland

^C Service de Neurochirurgie, Clinique du Millénaire, Montpellier, France

^D Service de Neurochirurgie, CHU de Nantes, Nantes, France

^E Nantes Université, INSERM, CRCI2NA, Nantes, France

^F Service d'Anesthésie Réanimation, Hôpital Beaujon, Assistance Publique
 Hôpitaux de Paris, Clichy, France

^G Université Paris Cité, INSERM, NeuroDiderot, Paris, France

^H Service de Médecine Physique et Réadaptation, CHU de Rouen, Rouen,
 France

^I Université de Rouen, Rouen, France

^J Service de Neurochirurgie Pédiatrique, Hôpital Femme Mère Enfant,
 Hospices Civils de Lyon, Bron, France

^K Institut de Biomécanique Humaine Georges Charpak, Arts et Métiers
 ParisTech et Université de Paris, Paris, France

* Corresponding author.

E-mail address: romain.manet@chu-lyon.fr (R. Manet).